

# Locally Convex Analysis in Differential Geometry

## Minicourse: Who is afraid of Infinite Dimensions?

Michael Heins

October 2025 – Lyon

## Table of Contents

|                                            |    |
|--------------------------------------------|----|
| Table of Contents                          | 1  |
| 1 Beyond Normed Spaces                     | 2  |
| 2 From Seminorms to Locally Convex Spaces  | 5  |
| 3 Projective and Injective Tensor Products | 18 |
| 4 Nuclearity                               | 29 |
| Bibliography                               | 35 |

## Instruction Manual

These lecture notes are based on the textbooks [7, 8, 10–12, 15, 16] as well as the encyclopaedic [6]. The lecture notes [17, 19, 20] have also been instrumental. Finally, the three further textbooks [1, 13, 14] should be mentioned, as they each contain some particularly nice presentations of particular topics. Moreover, these notes harbour many scattered ideas and, undoubtedly, numerous typographical (and by all likelihood also some mathematical) errors introduced by the author. Moreover, the moreover, many interesting and sometimes crucial facts have been left as exercises. On a first reading, most of them can be assumed as blackboxes or ignored altogether, but they are of course crucial for a deeper understanding. The unsure or otherwise confused reader should thus not be afraid to reach out to discuss. The main goal of these notes is to provide a gentle introduction to locally convex analysis for the working differential geometer, while also ultimately covering some advanced topics regarding tensor products and the arcane notion of nuclearity. As such, we assume a certain familiarity with notions from point set topology, but stay on the lighter side regarding functional analysis. Finally, the author's background results in a general philosophy of *reducing problems to inequalities*, whereas the puristic topological point of view takes a secondary role.

If you are still here: Why are you reading the instruction manual? What kind of dodgy lecture notes require a manual anyway? Who sold you these? Be careful out there. There are *analysts* about. One anyway.

# 1 Beyond Normed Spaces

Throughout the text, the principal assumption is that we are working with vector spaces over the field  $\mathbb{C}$  of complex numbers. Accordingly, linear mappings are complex linear and most mappings take values in the complex numbers. The real theory is largely analogous, and we indicate whenever it is not.

While normed spaces provide a fruitful and vast framework, they turn out to be insufficient to capture a number of natural phenomena within Differential Geometry and Analysis. In this preliminary section, we explore several such examples.

**Example 1.1** Let  $K \subseteq \mathbb{C}^n$  be compact. Then we may endow the space of complex-valued continuous functions

$$\mathcal{C}(K) := \{f: K \longrightarrow \mathbb{C} \mid f \text{ is continuous}\}$$

with the supremum norm

$$\|f\|_K := \max_{z \in K} |f(z)| \quad (1.1)$$

to obtain a Banach space. The corresponding topology of *uniform convergence* may be defined by taking the collection of open balls

$$B_r(f) := \{g \in \mathcal{C}(K) : \|f - g\|_K < r\}$$

with  $r > 0$  as a topological basis. That is to say, a subset  $\mathcal{U} \subseteq \mathcal{C}(K)$  is open iff it may be written as a union of open balls. The compactness of  $K$  is used to ensure that the maximum within (1.1) is well defined. Hence, it is a natural question how one should go about topologizing

$$\mathcal{C}(U) := \{f: U \longrightarrow \mathbb{C} \mid f \text{ is continuous}\}$$

for, say open, subsets  $U \subseteq \mathbb{C}^n$ . Two natural wishes are the continuity of the restriction mappings

$$\cdot|_K : \mathcal{C}(U) \longrightarrow \mathcal{C}(K) \quad (1.2)$$

for any compact set  $K \subseteq U$  and the completeness of the resulting space. Taking a step back, we remember that continuity is a local property, and as such not only preserved by uniform convergence, but also by *locally uniform convergence*. By virtue of local compactness of  $\mathbb{C}^d$ , this may be rephrased as uniform convergence on compact subsets. That is to say, a net within  $\mathcal{C}(U)$  should converge iff all of its images under the restriction mappings (1.2) are convergent in the normed spaces  $\mathcal{C}(K)$ . By choosing an exhaustion  $(K_n)_n$  of  $U$  by compact subsets, this may in turn be formalized by introducing the metric

$$d(f, g) := \sum_{n=0}^{\infty} \frac{1}{2^n} \cdot \frac{\|f - g\|_{K_n}}{1 + \|f - g\|_{K_n}} \quad \text{for all } f, g \in \mathcal{C}(U). \quad (1.3)$$

The resulting metric is invariant under translations and turns  $\mathcal{C}(U)$  into a complete metric space by Exercise 1.3. We have moreover built in the continuity of (1.2) into the definition. However, from a topological point of view, the open sets seem at first glance rather complicated. Taking a closer look, we realize that the “open cylinders”

$$B_{K,r}(f) := \{g \in \mathcal{C}(U) : \|f - g\|_K < r\} \subseteq \mathcal{C}(U) \quad (1.4)$$

are actually open for all compact subsets  $K \subseteq U$ ,  $r > 0$  and  $f \in \mathcal{C}(U)$ . Indeed, if  $g \in B_{K_n, r}(f)$  and  $h \in B_{2^{-n} \cdot r_0}(g)$  is in the metric ball with radius  $r_0 < 1$  around  $g$ , then using that

$$\phi: [0, \infty) \longrightarrow [0, 1), \quad \phi(x) := \frac{x}{1+x}$$

is strictly increasing with inverse  $\phi^{-1}(y) = y/(1-y)$  yields

$$\frac{\|g - h\|_{K_n}}{1 + \|g - h\|_{K_n}} < r_0 \quad \implies \quad \|g - h\|_{K_n} < \frac{r_0}{1 - r_0} = \phi^{-1}(r_0)$$

and thus

$$\|f - h\|_{K_n} \leq \|f - g\|_{K_n} + \|g - h\|_{K_n} < \|f - g\|_{K_n} + \phi^{-1}(r_0)$$

Hence, setting

$$r_0 := \phi(r - \|f - g\|_{K_n}) \in [0, 1)$$

produces a metric ball contained within  $B_{K, r}(f)$ , proving the openness. Conversely, every  $g \in B_r(f)$  is contained within  $B_{K, \phi^{-1}(2^{-n}r)}(f)$  for all  $n \in \mathbb{N}_0$  and thus we may forget about the metric balls in the sequel and only work with the open cylinders. Having established this, we get another pleasant property of this topology essentially for free. Namely, the continuity of the pointwise vector space operations

$$\begin{aligned} +: \mathcal{C}(U) \times \mathcal{C}(U) &\longrightarrow \mathcal{C}(U) \\ \cdot: \mathbb{C} \times \mathcal{C}(U) &\longrightarrow \mathcal{C}(U) \end{aligned}$$

and even of the pointwise multiplication. Geometrically, the sets (1.4) may indeed be thought of as cylinders, as it is absolutely convex and  $\|\cdot\|_K$  comes with the typically sizable kernel

$$\ker(\|\cdot\|_K) = \{f \in \mathcal{C}(U): f|_K \equiv 0\}.$$

Algebraically, the mappings  $\|\cdot\|_K$  thus constitute seminorms on  $\mathcal{C}(U)$ . That is to say, they fulfil all properties of a norm, except for having trivial kernel. By what we have shown, they may be used as a basis for the metric topology. This correspondence between systems of seminorms and topologies is at the heart of locally convex topologies and we shall explore this in Section 2.

It is instructive to work out the details of the arguments used in the prior discussion.

**Exercise 1.2** Let  $X$  be a metric space. Prove that a subset  $U \subseteq X$  is open iff it is a union of open balls.

**Exercise 1.3** Let  $X$  be a locally compact topological space and  $f, f_\alpha: X \longrightarrow \mathbb{C}$  be mappings for all  $\alpha \in J$  for some directed set  $J$ . Prove that the following are equivalent:

- i.) For every compact set  $K \subseteq X$ , the net of the restrictions  $(f_\alpha|_K)$  converges uniformly to the function  $f|_K$ .
- ii.) Every  $p \in X$  has an open neighbourhood  $U \subseteq X$  such that the net of restrictions  $(f_\alpha|_U)$  converges uniformly to the function  $f|_U$ .

**Exercise 1.4** Let  $U \subseteq \mathbb{R}^d$  or  $U \subseteq \mathbb{C}^n$  be open. Prove that the metric (1.3) turns  $\mathcal{C}(U)$  into a complete metric space.

Before turning to the general theory, we discuss some more remarkable observations that necessitate the investigation of functional analysis beyond norms.

**Example 1.5** Consider the space  $\mathcal{C}^\infty(\mathbb{R})$  of smooth functions on the real line and the differentiation operator

$$D := \frac{d}{dx}: \mathcal{C}^\infty(\mathbb{R}) \longrightarrow \mathcal{C}^\infty(\mathbb{R}).$$

Then the exponential functions

$$f_\alpha: \mathbb{R} \longrightarrow \mathbb{C}, \quad f_\alpha(x) := \exp(\alpha x)$$

constitute eigenfunctions of  $D$  for all  $\alpha \in \mathbb{C}$ . Hence, even if we somehow succeed in endowing  $\mathcal{C}^\infty(\mathbb{R})$  with the structure of a Banach algebra, then  $D$  is necessarily discontinuous by virtue of its unbounded spectrum.

Another strange feature of normed spaces is that some are never complete, regardless of how one chooses the norm.

**Lemma 1.6** *Let  $V$  be a normed vector space of countably infinite dimension. Then  $V$  is incomplete.*

PROOF: Assuming otherwise, we consider the family of closed subspaces

$$V_n := \text{span}\{e_1, \dots, e_n\} \quad \text{for } n \in \mathbb{N}_0$$

and where  $(e_n)_n \subseteq V$  is any basis. Then

$$V = \bigcup_{n=0}^{\infty} V_n,$$

and thus there exists some  $n_0 \in \mathbb{N}$  such that  $V_{n_0}^\circ \neq \emptyset$  by Baire's Theorem. This is a contradiction, as  $z e_{n_0+1} \notin V_{n_0}$  for all  $z \in \mathbb{C} \setminus \{0\}$ .  $\square$

Finally, we recast some classical complex analysis in a locally convex light.

**Exercise 1.7** Let  $U \subseteq \mathbb{C}^d$  be open. Prove that the space of holomorphic functions  $\mathcal{H}(U)$  constitutes a closed subspace of  $\mathcal{C}(U)$ .

Recall Montel's Theorem.

**Theorem 1.8 (Montel)** *Let  $U \subseteq \mathbb{C}^d$  be a domain. Then every locally bounded sequence  $(f_n)_n \subseteq \mathcal{H}(U)$  has a convergent subsequence.*

Classically, one speaks of normality of such families of functions. The reason for this is that Montel's work actually predates the abstract notion of compactness. Either way, we get the following consequence.

**Lemma 1.9** *Let  $U \subseteq \mathbb{C}^d$  be a domain. Then the topology of locally uniform convergence on the space of holomorphic functions  $\mathcal{H}(U)$  on  $U$  may not be induced by a norm.*

PROOF: Assuming otherwise, we get that the open unit ball

$$B_1(0) \subseteq \mathcal{H}(U)$$

constitutes a bounded subset of  $\mathcal{H}(U)$ . By Montel's Theorem, this implies the compactness of its closure  $B_1(0)^{\text{cl}}$ , which is only possible if  $\mathcal{H}(U)$  is finite-dimensional. This is a contradiction to the holomorphicity of the monomials.  $\square$

In this sense, the space  $\mathcal{H}(U)$  behaves much like a finite-dimensional space. The abstract underlying reason is its nuclearity, which is the content of Section 4.

## 2 From Seminorms to Locally Convex Spaces

Having convinced ourselves of the wonders and richness of the locally convex world within some examples, we now venture into the its abstract horrors. Most of the material is standard and can be found in all of the textbooks mentioned in the introduction. That being said, a particularly conceptually clean presentation is [11, Ch. 3]. As already noted, we shall place our focus on seminorms, as they ultimately frame the abstract results in a way convenient for the investigation of concrete problems.

**Definition 2.1 (Seminorms)** *Let  $V$  be a vector space. A mapping  $q: V \longrightarrow [0, \infty)$  is called a seminorm if:*

*i.) It is absolutely 1-homogeneous, i.e.*

$$q(\lambda \cdot v) = |\lambda| \cdot q(v) \quad \text{for all } v \in V, \lambda \in \mathbb{C}. \quad (2.1)$$

*ii.) It fulfils the triangle inequality*

$$q(v + w) \leq q(v) + q(w) \quad \text{for all } v, w \in V.$$

Note that (2.1) implies  $q(0) = 0$ , as we are working in characteristic different from two. Associated to any seminorm  $q$ , we define its collection of open cylinders

$$B_{q,r}(v) := \{w \in V : q(v - w) < r\}$$

with radius  $r > 0$ . We proceed with a list of examples.

**Example 2.2 (Seminorms)**

*i.) Every norm is a seminorm. The corresponding open cylinders are the open balls we are all used to.*

*ii.) Let  $v': V \longrightarrow \mathbb{C}$  be linear. Then*

$$|v'| := |\cdot| \circ v'$$

*is a seminorm of particularly simple type. An instructive example is the projection*

$$v': \mathbb{R}^2 \longrightarrow \mathbb{C}, \quad v'(x^1, x^2) := x^1$$

onto the first component. Then

$$\ker|v'| = \ker v' = \{0\} \times \mathbb{R}$$

as well as

$$|v'|((x^1, x^2) - (y^1, y^2)) = |x^1 - y^1| \quad \text{for all } x, y \in \mathbb{R}^2.$$

Consequently,

$$B_{|v'|,r}(x) = \{y \in \mathbb{R}^2 : |x^1 - y^1| < r\} = (x^1 - r, x^1 + r) \times \mathbb{R}$$

really constitutes a cylinder in this case.

iii.) Let  $U \subseteq \mathbb{C}^d$  be open. Then mappings  $\|\cdot\|_K$  constitute seminorms on  $\mathcal{C}(U)$  for all compact subsets  $K \subseteq U$ . More generally, we get seminorms on  $\mathcal{H}(U)$  by setting

$$\|f\|_{K,\alpha} := \max_{|\beta| \leq |\alpha|} \|\partial^\beta f\|_K$$

for any multi-index  $\alpha \in \mathbb{N}_0^d$ . Replacing complex derivatives with real ones, the same formula also defines seminorms on  $\mathcal{C}^\infty(U)$ . By virtue of the identity principle, if  $U$  is connected, then  $\|\cdot\|_K$  is even a norm on  $\mathcal{H}(U)$  for every compact  $K \subseteq U$  with non-empty open interior.

iv.) For every  $r > 0$ , the mappings

$$q_r \left( \sum_{n=0}^{\infty} a_n \cdot z^n \right) := \sum_{n=0}^{\infty} |a_n| \cdot r^n$$

define norms on  $\mathcal{H}(\mathbb{C})$ .

v.) Let  $n, m \in \mathbb{N}_0$  and define

$$r_{n,m}(f) := \sup_{x \in \mathbb{R}} (1 + x^2)^m \cdot |f^n(x)|$$

for  $f \in \mathcal{C}^\infty(\mathbb{R})$ . Then  $r_{n,m}(f) < \infty$  for all  $n, m \in \mathbb{N}_0$  is equivalent to  $f$  being an element of the classical Schwartz space. Once again, we are actually dealing with norms instead of just seminorms.

It is a common theme within locally convex analysis to rephrase the membership in some function space as a seminorm condition. As we shall see, this then gives rise to an associated locally convex topology, which often turns out to have desirable properties such as completeness.

**Exercise 2.3** Show that the systems of seminorms

$$\{\|\cdot\|_K : K \subseteq \mathbb{C} \text{ compact}\}$$

and

$$\{q_r : r > 0\}$$

are equivalent on  $\mathcal{H}(\mathbb{C})$ . That is to say, for every compact subset  $K \subseteq \mathbb{C}$  there exists some  $r > 0$  with

$$\|f\|_K \leq q_r(f) \quad \text{for all } f \in \mathcal{H}(\mathbb{C})$$

and, conversely, for every  $r > 0$  there exists a compact set  $K \subseteq \mathbb{C}^d$  such that

$$q_r(f) \leq \|f\|_K \quad \text{for all } f \in \mathcal{H}(\mathbb{C}).$$

Remarkably, one may reconstruct a seminorm from its open unit cylinder. This correspondence between algebraic gadgets and geometric objects is established by means of the Minkowski functional.

**Definition 2.4 (Minkowski functional)** *Let  $V$  be a vector space and  $U \subseteq V$  be absorbing, i.e.*

$$\bigcup_{r>0} rU = \bigcup_{r>0} \{r \cdot v : v \in U\} = V.$$

*Then*

$$p_U : V \longrightarrow [0, \infty), \quad p_U(v) := \inf \{r > 0 : v \in r \cdot U\} \quad (2.2)$$

*is called the Minkowski functional of  $U$ .*

Recall that a subset  $C$  of a vector space  $V$  is called convex if the line segment

$$[v, w] := \{\lambda \cdot v + (1 - \lambda)w : \lambda \in [0, 1]\} \subseteq C$$

for all  $v, w \in C$ . It is called balanced (or cyclic) if

$$\eta \cdot U = U \quad \text{for all } \eta \in \mathbb{C}, |\eta| = 1.$$

Combining both notions, one speaks of absolute convexity of  $U$ .

**Proposition 2.5** *Let  $V$  be a vector space and  $U \subseteq V$  be absorbing.*

*i.) The Minkowski functional (2.2) is well-defined.*

*ii.) If  $U$  is convex, then  $p_U$  is sublinear, i.e.*

$$p_U(v + w) \leq p_U(v) + p_U(w) \quad \text{and} \quad p(r \cdot v) = r \cdot p(v)$$

*for all  $v, w \in V$  and  $r \geq 0$ .*

*iii.) If  $U$  is convex and balanced, then  $p_U$  is a seminorm. Moreover,*

$$B_{p_U,1}(0) \subseteq U \subseteq B_{p_U,1}(0)^{\text{cl}}.$$

*iv.) If  $q$  is a seminorm, then the open cylinders  $B_{q,r}(0)$  are absorbing, convex and balanced for all  $r > 0$ . Moreover,*

$$p_{B_{q,1}(0)} = q = p_{B_{q,1}(0)^{\text{cl}}},$$

*where we define*

$$B_{q,1}(0)^{\text{cl}} = \{v \in V : q \leq 1\}.$$

**Exercise 2.6** Prove Proposition 2.5.

The central idea behind locally convex topologies now to use the open cylinders corresponding to a collection of seminorms as the basis of a topology. This is also where the phrase *locally convex* stems from: The origin possesses a neighbourhood basis consisting of absolutely convex sets! By Proposition 2.5, this demand automatically leads towards seminorms.

Returning to this idea, we define a topology on  $V$  associated to some collection  $\mathcal{P}$  of seminorms by first taking finite intersections of open cylinders and then arbitrary unions of the resulting sets. There is a simple condition on the level of seminorms, as to when we may skip the first step, i.e. are actually dealing with a basis of the topology. Note that any set of seminorms is, in particular, a set of real-valued mappings defined on a joint domain. As such, it carries the pointwisely defined partial order, i.e.

$$q \leq p \iff q(v) \leq p(v) \text{ for all } v \in V.$$

Recall that a partially ordered set  $(\mathcal{P}, \leq)$  is called a directed if, i.e. for all  $q_1, q_2 \in \mathcal{P}$  there exists  $p \in \mathcal{P}$  with  $q_1 \leq p$  and  $q_2 \leq p$ .

**Lemma 2.7** *Let  $V$  be a vector space and  $\mathcal{P}$  a set of seminorms on  $V$  such that  $(\mathcal{P}, \leq)$  is a directed set. Then every finite intersection of elements within the set*

$$\mathcal{B} := \{B_{q,r}(v) : q \in \mathcal{P}, r > 0, v \in V\} \quad (2.3)$$

*may be written as a union of elements of  $\mathcal{B}$ .*

PROOF: It suffices to prove the claim for intersections of two open cylinders. To this end, let  $q_1, q_2 \in \mathcal{P}$ ,  $r_1, r_2 > 0$  and  $v_1, v_2 \in V$  such that

$$U := B_{q_1, r_1}(v_1) \cap B_{q_2, r_2}(v_2) \neq \emptyset.$$

By assumption, we find a joint upper bound  $p \in \mathcal{P}$  of  $q_1$  and  $q_2$ . Let  $v \in U$  and

$$r_v := \min\{r_1 - q_1(v - v_1), r_2 - q_2(v - v_2)\} > 0.$$

Then  $B_{p, r_v}(v) \subseteq U$ , as

$$q_1(v - w) \leq q_1(v - v_1) + p(v_1 - w) < q_1(v - v_1) + r_v < r_1$$

and, analogously,  $q_2(v - w) < r_2$  for all  $w \in B_{p, r_v}(v)$ . Consequently, we get

$$U = \bigcup_{v \in U} B_{p, r_v}(v),$$

as  $v \in B_{p, r_v}(v)$  for all  $v \in U$ . □

It is customary to call a collection of seminorms  $\mathcal{P}$  *filtrating* if  $(\mathcal{P}, \leq)$  is directed. As linear combinations of seminorms with non-negative coefficients are again seminorms, the collection of all seminorms is always filtrating. We are now in a position to define locally convex spaces.

**Definition 2.8 (Locally convex space)** *A vector space  $V$  endowed with a topology is called locally convex if there exists a filtrating system of seminorms  $\mathcal{P}$  on  $V$  such that*

$$\mathcal{B} := \{B_{q,r}(v) : q \in \mathcal{P}, r > 0, v \in V\}$$

*constitutes a basis of the topology. We then call  $\mathcal{P}$  a defining system of seminorms for  $V$ . Moreover, we write*

$$\text{cs}(V) := \{q : V \longrightarrow \mathbb{R} \mid q \text{ is a continuous seminorm}\}$$

*for the set of all continuous seminorms of a locally convex space.*

This definition raises some questions, the answers of which we collect before proceeding.

**Proposition 2.9** *Let  $V$  be locally convex with defining system of seminorms  $\mathcal{P}$ .*

*i.) The locally convex space  $V$  is a topological vector space, i.e.*

$$+: V \times V \longrightarrow V \quad \text{and} \quad \cdot: \mathbb{C} \times V \longrightarrow V$$

*are continuous, where we endow  $V \times V$  and  $\mathbb{C} \times V$  with the product topology.*

*ii.) For every  $v \in V$ , the translation*

$$\tau_v: V \longrightarrow V, \quad \tau_v(w) := v + w$$

*constitutes a linear homeomorphism.*

*iii.) A seminorm  $q$  on  $V$  is continuous iff there exist  $p_1, \dots, p_n \in \mathcal{P}$  and  $c_1, \dots, c_n > 0$  such that*

$$q \leq c_1 \cdot p_1 + \dots + c_n \cdot p_n. \quad (2.4)$$

*iv.) The locally convex topology induced by the collection of all continuous seminorm coincides with the topology of  $V$ .*

*v.) A subset  $U \subseteq V$  is open iff for every  $v \in U$  there exists  $q \in \text{cs}(V)$  and some  $r > 0$  with*

$$B_{q,r}(v) \subseteq U.$$

*vi.) The locally convex space  $V$  is Hausdorff iff for every  $v \in V$  there exists  $q \in \text{cs}(V)$  with  $q(v) > 0$ .*

PROOF: Let  $q \in \mathcal{P}$ ,  $r > 0$  and  $v \in V$ . We have to prove that

$$+^{-1}(B_{q,r}(v)) = \{(v_1, v_2) \in V \times V : q(v_1 + v_2 - v) < r\}$$

is open. Let  $(v_1, v_2) \in +^{-1}(B_{q,r}(v))$  and set  $\delta := q(v_1 + v_2 - v)$ . We claim

$$B := B_{q,(r-\delta)/2}(v_1) \times B_{q,(r-\delta)/2}(v_2) \subseteq +^{-1}(B_{q,r}(v)).$$

Indeed, if  $(w_1, w_2) \in B$ , then by the triangle inequality

$$q(w_1 + w_2 - v) \leq q(w_1 - v_1) + q(w_2 - v_2) + q(v_1 + v_2 - v) < \frac{r-\delta}{2} + \frac{r-\delta}{2} + \delta = r.$$

As  $B \subseteq V \times V$  is open in the product topology as a product of open sets, this implies the continuity of the vector space addition. Similarly, for the multiplication by scalars, we construct open neighbourhoods of

$$(z_0, w_0) \in \cdot^{-1}(B_{q,r}(v)) = \{(z, w) \in \mathbb{C} \times V : q(zw - v) < r\}.$$

Indeed, if  $(z, w) \in B_{r_1}(z_0) \times B_{q,r_2}(w_0)$  with  $\delta := q(z_0 w_0 - v)$ , then

$$\begin{aligned} q(zw - v) &\leq q(zw - z_0 w_0) + q(z_0 w_0 - v) \\ &= |z_0| \cdot q(w - w_0) + |z - z_0| \cdot q(w_0) + q(z_0 w_0 - v) \\ &< |z_0| \cdot r_2 + r_1 \cdot q(w_0) + \delta. \end{aligned}$$

Hence, setting

$$r_1 := r_2 := \frac{r - \delta}{2(1 + q(w_0) + |z_0|)}$$

implies

$$B_{r_1}(z_0) \times B_{q,r_2}(w_0) \subseteq \cdot^{-1}(B_{q,r}(v)).$$

We have thus shown *i.*). Now, *ii.*) is clear, as the inclusions  $V \hookrightarrow V \times V$  are topological embeddings. For *iii.*), let  $q \in \text{cs}(V)$ . Then, by definition of continuity, the preimage

$$q^{-1}([0, 1)) \subseteq V$$

is open. Hence, it can be written as a union of open cylinders corresponding to elements of  $\mathcal{P}$ . In particular, we find  $p \in \mathcal{P}$  and  $r > 0$  such that

$$B_{p,r}(0) \subseteq q^{-1}([0, 1)) = B_{q,1}(0),$$

where we use  $q(0) = 0$ .<sup>1</sup> Unwrapping the definition of the open cylinder, this means

$$p(v) < r \quad \implies \quad q(v) < 1$$

for all  $v \in V$ . By homogeneity, we may rephrase this as the continuity estimate

$$q(v) \leq r \cdot p(v) \quad \text{for all } v \in V.$$

This is (2.4), where we may thus even choose  $n = 1$ . Assume conversely (2.4) holds for some seminorm  $q$  and suitably chosen  $p_1, \dots, p_n \in \mathcal{P}$  and  $c_1, \dots, c_n > 0$ . As  $\mathcal{P}$  is filtrating, we may pass to a joint upper bound  $p \in \mathcal{P}$  of  $p_1, \dots, p_n$  to obtain

$$q \leq c \cdot p$$

with  $c := c_1 + \dots + c_n$ . Reversing the logic from before, this means

$$B_{p,1}(0) \subseteq B_{q,1/c}(0) = q^{-1}([0, 1/c)).$$

Hence,  $p$  is continuous at the origin. Let now  $(v_\alpha)_{\alpha \in J}$  be a convergent net with limit  $v \in V$ . Then the net  $(v - v_\alpha)_\alpha$  converges to zero by *ii.*). By what we have already shown, given  $\varepsilon > 0$  we thus find  $\alpha_0 \in J$  such that

$$p(v_\alpha) \leq p(v) + p(v - v_\alpha) \leq \varepsilon \quad \text{for all } \alpha \succ \alpha_0.$$

Bringing  $p(v)$  to the other side and varying  $\varepsilon$  implies  $p(v_\alpha) \rightarrow p(v)$ , establishing the continuity of  $p$  on all of  $V$ . This completes the proof of *iii.*). We turn towards *iv.*). By Definition 2.8, it is clear that the topology induced by  $\text{cs}(V)$  is finer than the one induced by  $\mathcal{P}$ . It thus suffices to prove that given  $q \in \text{cs}(V)$ ,  $r > 0$  and  $v \in V$ , the associated open cylinder  $B_{q,r}(v)$  is already a member of the topology generated by  $\mathcal{P}$ . For  $v = 0$ , this is just the geometric formulation of *iii.*). Invoking *ii.*), this implies the openness of

$$\tau_v(B_{q,r}(0)) = B_{q,r}(v)$$

as homeomorphisms are, in particular, open mappings. Part *v.*) is now immediate, as each  $B_{q,r}(v)$  for  $q \in \text{cs}(V)$  contains an open cylinder associated to a seminorm from the

---

<sup>1</sup>If you are not familiar with this type of reasoning within topological spaces, prove this!

defining system  $\mathcal{P}$  by what we have just shown. Finally, assume the Hausdorff property. Then, we may separate  $v \neq 0$  from the origin. In particular, there exist  $q \in \mathcal{P}$  and  $r > 0$  with

$$v \notin B_{q,r}(0),$$

which means  $q(v) > r > 0$ . Conversely, assume the condition and let  $v, w \in V$  with  $v \neq w$ . Then  $v - w \neq 0$  and thus we may invoke our assumption to find  $q \in \text{cs}(V)$  such that  $r := q(v - w) > 0$ . Consequently,

$$B_{q,r/2}(0) \cap B_{q,r/2}(v - w) = \emptyset$$

by the triangle inequality. Translating by  $w$  via *ii.*) then yields

$$B_{q,r/2}(w) \cap B_{q,r/2}(v) = \emptyset.$$

Hence,  $V$  is Hausdorff in this case.  $\square$

In the proof of *iii.*), we have seen that it suffices to consider the case  $n = 1$ . Our more general formulation is however often useful in practice. The following exercise ties a loose end from earlier.

**Exercise 2.10** Let  $V$  be locally convex,  $q \in \text{cs}(V)$  and  $r > 0$ . Prove that the open cylinder  $B_{q,r}(v)$  is sequentially dense in the closed cylinder

$$B_{q,r}(v)^{\text{cl}} := \{w \in V : q(v - w) \leq r\}.$$

This, a posteriori, justifies our notation.

Next, we take a brief look at how convergence within a locally convex space may be described. Ultimately, the generalization is both straightforward and convenient.

**Lemma 2.11** *Let  $V$  be a locally convex space with defining system of seminorms  $\mathcal{P}$ . Then a net  $(v_\alpha)_{\alpha \in J}$  converges to a vector  $v \in V$  iff for every  $q \in \mathcal{P}$  and every  $\varepsilon > 0$ , there exists an index  $\alpha_0 \in J$  such that*

$$q(v_\alpha - v) \leq \varepsilon \quad \text{for all } \alpha \succ \alpha_0. \quad (2.5)$$

*In this case, the condition holds for all  $q \in \text{cs}(V)$ .*

PROOF: Assume first that the net  $(v_\alpha)_{\alpha \in J}$  is convergent with limit  $v \in V$  and let  $q \in \text{cs}(V)$ . Then the open cylinder  $B_{q,\varepsilon}(v)$  constitutes an open neighbourhood of  $v$  by Proposition 2.9, *iv.*) and *v.*). By convergence of  $(v_\alpha)_{\alpha \in J}$  there thus is some  $\alpha_0 \in J$  such that  $v_\alpha \in B_{q,\varepsilon}(v)$  for all  $\alpha \succ \alpha_0$ . This is precisely (2.5). Conversely, let again  $\varepsilon > 0$  as well as  $q \in \mathcal{P}$  and assume the validity of (2.5). Then  $v_\alpha \in B_{q,\varepsilon}(v)$  for all  $\alpha \succ \alpha_0$ . As the open cylinders corresponding to seminorms from the defining system form a basis of the topology, this already proves the convergence of  $(v_\alpha)$  to  $v$ .  $\square$

Motivated by the Lemma, we call a net  $(v_\alpha)_{\alpha \in J} \subseteq V$  within a locally convex space  $V$  Cauchy if for every  $q \in \mathcal{P}$  and every  $\varepsilon > 0$  there exists an index  $\alpha_0 \in J$  such that

$$q(v_\alpha - v_\beta) \leq \varepsilon \quad \text{for all } \alpha, \beta \succ \alpha_0.$$

In this case, the condition holds for all  $q \in \text{cs}(V)$ . As usual, we then call  $V$  complete if all Cauchy nets converge. If the topology of  $V$  actually arises from a single norm, then this indeed recovers the usual notions of Cauchy nets and, by first countability, of completeness.

**Exercise 2.12** Let  $V$  be a vector space and  $\|\cdot\|$  a norm on  $V$ . Prove that the locally convex topology induced by  $\mathcal{P} := \{\|\cdot\|\}$  is the norm-topology. Can you describe the set of all continuous seminorms  $\text{cs}(V)$  on  $V$ ?

Each of the systems of seminorms from Example 2.2 now induces the corresponding vector spaces with locally convex topologies.

**Exercise 2.13** Consider the space of complex sequences

$$\text{Map}(\mathbb{N}_0, \mathbb{C}) := \{a: \mathbb{N}_0 \longrightarrow \mathbb{C}\}.$$

i.) Construct a defining system of seminorms such that the resulting locally convex topology is the topology of pointwise convergence.

Hint: Remember that defining systems are filtrating.

ii.) Establish the completeness of the locally convex space  $\text{Map}(\mathbb{N}_0, \mathbb{C})$ .

iii.) Let  $\|\cdot\|$  be a norm on  $\text{Map}(\mathbb{N}_0, \mathbb{C})$ . Prove that  $\|\cdot\|$  is discontinuous.

iv.) Show that the space

$$c_{00} := \{a \in \text{Map}(\mathbb{N}_0, \mathbb{C}) : \exists N \in \mathbb{N}_0 \forall n \geq N \ a_n = 0\}.$$

of compactly supported sequences is dense within  $\text{Map}(\mathbb{N}_0, \mathbb{C})$ .

Hint: It is *not* sequentially dense. Use nets.

**Proposition 2.14** Let  $L: V \longrightarrow W$  be a linear mapping between locally convex spaces  $V$  and  $W$  with corresponding defining systems of seminorms  $\mathcal{P}_V$  and  $\mathcal{P}_W$ . Then the following are equivalent:

- i.) The map  $L$  is uniformly continuous.
- ii.) The map  $L$  is continuous.
- iii.) The map  $L$  is continuous at some  $v \in V$ .
- iv.) The map  $L$  is continuous at the origin.
- v.) For every  $q \in \text{cs}(V)$  there exist  $p \in \mathcal{P}_V$  and  $c > 0$  such that

$$q(Lv) \leq c \cdot p(v) \quad \text{for all } v \in V. \quad (2.6)$$

vi.) For every  $q \in \mathcal{P}_W$  there exists  $p \in \text{cs}(V)$  such that

$$q(Lv) \leq p(v) \quad \text{for all } v \in V.$$

PROOF: Clearly, i.) implies ii.), which in turn implies iii.). Assuming  $L$  is continuous at a point  $v \in V$  implies the continuity of the composition

$$w \mapsto (\tau_{-Lv} \circ L \circ \tau_v)(w) = L(v) - L(v + w) = -Lw$$

at  $w = 0$ , as  $\tau_v(0) = v$  and by Proposition 2.9, ii.). This is iv.). Alternatively, this follows readily by checking the continuity at zero by means of nets and using the linearity. Assume now iv.) and let  $q \in \text{cs}(V)$ . As  $B_{q,1}(0)$  constitutes an open neighbourhood of zero, the continuity yields the openness of its preimage under  $L$ . As the open cylinders

centered at zero corresponding to  $\mathcal{P}_V$  form a neighbourhood basis, we thus find  $p \in \mathcal{P}_V$  and  $c > 0$  such that

$$B_{p,1/c}(0) \subseteq L^{-1}(B_{q,1}(0)).$$

Unwrapping this inclusion yields precisely (2.6). As multiples of continuous seminorms are continuous,  $v.)$  implies  $vi.)$ . Finally, assume  $vi.)$  and let  $q \in \mathcal{P}_W$  as well as  $r > 0$ . By rescaling our assumption, we find  $p \in cs(V)$  such that

$$q(Lv) \leq p(v) \quad \text{for all } v \in V.$$

That is to say,

$$Lv - Lw = L(v - w) \in B_{q,r}(0) \quad \text{for all } v, w \in V \text{ such that } v - w \in B_{p,r}(0).$$

This is precisely the uniform continuity of  $L$ .  $\square$

Both  $v.)$  and  $vi.)$  are useful in practice: The former for using, the latter for checking continuity. It is instructive to apply Proposition 2.14 to some simple examples. They also illustrate the usefulness of working with small defining systems, while also being aware of many continuous seminorms.

**Exercise 2.15** Let  $V$  be a finite dimensional locally convex Hausdorff space. Prove that, choosing a basis  $(e_1, \dots, e_d)$  of  $V$ , the coordinate mapping

$$\Phi: \mathbb{C}^d \longrightarrow V, \quad \Phi(z^1, \dots, z^d) := \sum_{n=1}^d z^n e_n$$

constitutes a linear homeomorphism.

**Exercise 2.16** Prove the continuity of the differentiation operator

$$D: \mathcal{H}(\mathbb{C}) \longrightarrow \mathcal{H}(\mathbb{C}), \quad Df := f'$$

and the multiplication operator

$$M: \mathcal{H}(\mathbb{C}) \longrightarrow \mathcal{H}(\mathbb{C}), \quad (Mf)(z) := zf(z).$$

Compute also the commutator

$$[D, M] := D \circ M - M \circ D.$$

Conclude again that the topology of  $\mathcal{H}(\mathbb{C})$  can not be induced by a norm.

Hint: It might be convenient to take another look at Exercise 2.3 in light of the new technology we have established in the meantime.

The situation, in which we find a countable defining system of seminorms deserves particular attention. In this case, the locally convex topology is first countable, i.e. every point possesses a countable basis of neighbourhoods.

**Proposition 2.17 (Metrizability)** *Let  $V$  be a locally convex space. Then the following are equivalent:*

- i.) The space  $V$  is first countable.
- ii.) There exists a countable defining system of seminorms for  $V$ .
- iii.) There exists an ascending sequence  $(q_n)$  of seminorms defining the topology of  $V$ .
- iv.) The topology of  $V$  is metrizable by means of a translation invariant metric  $d$ , i.e.

$$d(v+x, w+x) = d(v, w) \quad \text{for all } v, w, x \in V.$$

- v.) The topology of  $V$  is metrizable.

PROOF: Assuming ii.), there exists a countable neighbourhood basis  $(U_n)_n$  of the origin. As the open cylinders centered at 0 corresponding to continuous seminorms on  $V$  constitute a neighbourhood basis themselves, we find corresponding seminorms  $p_n \in \text{cs}(V)$  such that

$$B_{p_n,1}(0) \subseteq U_n \quad \text{for all } n \in \mathbb{N}_0.$$

Translating these balls by means of Proposition 2.9, ii.) we get countable neighbourhood bases consisting of open cylinders corresponding to the  $p_n$ . Thus,  $\mathcal{P} := \{p_n : n \in \mathbb{N}_0\}$  constitutes a countable defining system of seminorms for the locally convex space  $V$ . As  $\text{cs}(V)$  is closed under taking pointwise maxima of finitely many elements, we may define

$$q_n := \max\{p_1, \dots, p_n\} \quad \text{for all } n \in \mathbb{N}_0$$

to obtain an ascending countable defining system of seminorms for  $V$ . Mimicking the construction from Example 1.1, this in turn yields a translation invariant metric  $d$  on  $V$  defined by

$$d(v, w) := \sum_{n=0}^{\infty} 2^{-n} \cdot \frac{q_n(v-w)}{1 + q_n(v-w)}.$$

Repeating our considerations regarding the metric balls establishes that  $d$  generates the topology of  $V$ . Finally, metrizable spaces are certainly first countable.  $\square$

If one (and thus all) of the conditions within Proposition 2.17 are fulfilled, and  $V$  is moreover complete Hausdorff, then we call  $V$  a Fréchet space. Banach spaces are particular examples of Fréchet spaces by Exercise 2.12. As the First Baire Category Theorem applies to complete metric spaces, it should be of no surprise that there are Fréchet space generalizations of the Banach-Steinhaus Theorem, the Open Mapping Theorem and the Closed Graph Theorem. In fact, appropriately phrased, they remain valid beyond the setting of Fréchet spaces. The exploration of these topics however goes way beyond the scope of these lecture notes. Discussions can be found in any of the excellent textbooks mentioned at the beginning of the text, where we once again highlight [11, Ch. 4 & 5] for a particularly pleasant experience.

Instead, we proceed with a locally convex incarnation of the Hahn-Banach Extension Theorem, which we shall need for our exploration of tensor products.

**Theorem 2.18 (Hahn-Banach)** *Let  $V$  be a vector space and  $q: V \rightarrow \mathbb{R}$  be sublinear. Furthermore, let  $U \subseteq V$  be a subspace and*

$$u': U \rightarrow \mathbb{C}$$

be a linear functional on  $U$  such that  $|u'| \leq q$ . Then there exists a linear functional  $v': V \rightarrow \mathbb{C}$  such that

$$v'|_U = u' \quad \text{and} \quad |v'| \leq q.$$

In view of Proposition 2.14, this means that we may extend continuous linear functional from subspaces in manner that preserves continuity estimates. This statement is ultimately a purely algebraic application of Zorn's Lemma. It is quite likely that you have already seen a sufficiently general incarnation before. Thus, instead of rehearsing the details, we conclude our abstract considerations with a quite useful consequence for later.

**Exercise 2.19** Let  $V$  be locally convex and  $q \in \text{cs}(V)$ . Then

$$q(v) = \sup_{|v'| \leq q} |v'(v)| \quad \text{for all } v \in V. \quad (2.7)$$

Another amusing application is the existence of so-called Banach limits.

**Exercise 2.20 (Banach Limits)** Let  $L: \ell^\infty(\mathbb{R}) \rightarrow \ell^\infty(\mathbb{R})$  be the left-shift on the space of real bounded sequences, i.e.  $(La)_n := a_{n+1}$  for all  $a \in \ell^\infty(\mathbb{R})$ . Prove that there exists a continuous linear functional  $\Lambda$  on  $\ell^\infty$  such that

$$\Lambda(La) = \Lambda(a) \quad \text{and} \quad \liminf_{n \rightarrow \infty} a_n \leq \Lambda(a) \leq \limsup_{n \rightarrow \infty} a_n$$

for all  $a \in \ell^\infty(\mathbb{R})$ .

Hint: Consider the functionals

$$\Lambda_n(a) := \frac{1}{n} \sum_{k=0}^{n-1} a_k$$

defined on the subspace

$$Y := \{a \in \ell^\infty(\mathbb{R}) : \lim_{n \rightarrow \infty} \Lambda_n(a) \text{ exists}\}.$$

After our abstract considerations on locally convex spaces, our goal is now to endow the space of complex-valued smooth functions  $\mathcal{C}^\infty(M)$  defined on a manifold  $M$  with a locally convex topology. In Example 2.2, *iii.*, we have already met seminorms for smooth functions defined on open subsets of  $\mathbb{R}^n$  or  $\mathbb{C}^n$ . As usual in differential geometry, the idea is then that this reflects the situation within a chart. While this approach is perfectly serviceable, there is a more conceptual (and ultimately equivalent) approach based on differential operators and having already established the  $\mathcal{C}$ -topology, see again Example 1.1.

**Definition 2.21 ( $\mathcal{C}$ -topology)** Let  $M$  be a topological space. Then the  $\mathcal{C}$ -topology on the space of complex-valued continuous functions  $\mathcal{C}(M)$  is the locally convex topology generated by the seminorms

$$q_K(f) := \max_{p \in K} |f(p)|,$$

where we vary  $K \subseteq M$  through the compact subsets of  $M$ .

As Recall Grothendieck's [4] recursive definition of differential operators of an associative algebra  $\mathcal{A}$  as it is e.g. discussed in [9, Ch. 15]. That is,  $\text{DiffOp}(\mathcal{A})$  is the filtered algebra

$$\begin{aligned}\text{DiffOp}^0(\mathcal{A}) &:= \{M_a : a \in \mathcal{A}\}, \\ \text{DiffOp}^k(\mathcal{A}) &:= \{D \in \text{L}(\mathcal{A}) \mid \forall_{a \in \mathcal{A}} : [D, M_a] \in \text{DiffOp}^{k-1}(\mathcal{A})\} \quad \text{for } k > 0,\end{aligned}$$

where  $\text{L}(\mathcal{A})$  is the space of  $\mathbb{k}$ -linear maps from  $\mathcal{A}$  to  $\mathcal{A}$ , the  $M_a$  are multiplication operators with  $a \in \mathcal{A}$  and

$$[\cdot, \cdot] : \text{L}(\mathcal{A}) \times \text{L}(\mathcal{A}) \longrightarrow \text{L}(\mathcal{A}), \quad [D, D'] := D \circ D' - D' \circ D$$

is the commutator induced from the associative algebra structure of  $\text{L}(\mathcal{A})$ . If  $\mathcal{A}$  is unital, then

$$\text{DiffOp}^0(\mathcal{A}) \cong \text{End}_{\mathcal{A}}(\mathcal{A}) \cong \mathcal{A},$$

where  $\text{End}_{\mathcal{A}}(\mathcal{A})$  denotes the set of  $\mathcal{A}$ -linear endomorphisms of  $\mathcal{A}$ . The algebra we are interested in is of course  $\mathcal{A} = \mathcal{C}^k(M)$  with  $k \in \mathbb{N} \cup \{\infty\}$ .

**Exercise 2.22** Let  $M$  be a manifold and  $k \in \mathbb{N}$ . Prove

$$\text{DiffOp}^\ell(\mathcal{C}^k(M)) = \text{DiffOp}^k(\mathcal{C}^\infty(M)) \quad \text{for } \ell = 0, \dots, k.$$

In the sequel, we thus simply speak of  $\text{DiffOp}^\ell(M)$ .

More generally, one may consider  $\mathbb{C}$ -linear mappings

$$D : \mathcal{E} \longrightarrow \mathcal{F}$$

between unital  $\mathcal{A}$ -modules  $\mathcal{E}$  and  $\mathcal{F}$ . Accordingly, the multiplication operators  $M_a$  with  $a \in \mathcal{A}$  now act on  $\mathcal{E}$  and  $\mathcal{F}$  by module multiplication. Hence, we may also speak of differential operators

$$D : \Gamma^\infty(E) \longrightarrow \Gamma^\infty(F)$$

between spaces of sections of vector bundles  $\text{pr}_E : E \longrightarrow M$  and  $\text{pr}_F : F \longrightarrow M$  over the same base manifold  $M$ . With these preliminaries, we may define the  $\mathcal{C}^k$ -topologies for finite  $k$ . The idea is that we may view a differential operator of order  $k$  as a linear mapping

$$D : \mathcal{C}^k(M) \longrightarrow \mathcal{C}(M). \tag{2.8}$$

Whatever the  $\mathcal{C}^k$ -topology may entail, we certainly want all such mappings to be continuous. As we are already in possession of a topology on  $\mathcal{C}(M)$ , the natural choice thus becomes the initial topology, i.e. the coarsest topology such that all mappings (2.8) become continuous.

**Definition 2.23 ( $\mathcal{C}^k$ -topology)** *Let  $M$  be a manifold and  $k \in \mathbb{N}$ . The  $\mathcal{C}^k$ -topology on the space of  $k$ -times differentiable functions  $\mathcal{C}^k(M)$  is the initial topology with respect to the mappings*

$$D : \mathcal{C}^k(M) \longrightarrow \mathcal{C}(M),$$

where we vary  $D \in \text{DiffOp}^k(M)$ .

The definition for the  $\Gamma^k$ -topology of  $k$ -times differentiable sections is mutatis mutandis identical. Remarkably, initial topologies induced by linear mappings into locally convex spaces are automatically locally convex.

**Proposition 2.24** *Let*

$$L_\alpha: V \longrightarrow V_\alpha$$

*be linear mappings from a vector space  $V$  with values in locally convex spaces  $V_\alpha$ . Then the associated initial topology on  $V$  is locally convex. More precisely, if  $\mathcal{P}_\alpha$  are defining systems of seminorms for  $V_\alpha$  for all  $\alpha \in J$ , then*

$$\mathcal{P} := \{L_\alpha^* p_\alpha := p_\alpha \circ L_\alpha : \alpha \in J, p_\alpha \in \mathcal{P}_\alpha\} \quad (2.9)$$

*constitutes a defining system of seminorms for  $V$ .*

PROOF: By linearity of the  $\phi_\alpha$ , the pullbacks  $\phi_\alpha^* p_\alpha$  are indeed seminorms on  $V$ . As locally convex topologies are defined by means of the bases (2.3), we get a basis for the topology of  $V$ , which is given by

$$\mathcal{B} := \{L_\alpha^{-1}(B_{p_\alpha, r}(v_\alpha)) : \alpha \in J, p_\alpha \in \mathcal{P}_\alpha, r > 0, v_\alpha \in V_\alpha\}.$$

Notice now

$$\phi_\alpha^{-1}(B_{p_\alpha, r}(L_\alpha v)) = \{w \in V : p_\alpha(L_\alpha v - L_\alpha w) < r\} = B_{\phi_\alpha^* p_\alpha, r}(v)$$

for all  $w \in V$ ,  $\alpha \in J$  and  $r > 0$ . Now, if  $v_\alpha \notin L_\alpha V$ , then

$$\phi_\alpha^{-1}(B_{p_\alpha, r}(L_\alpha v)) = \emptyset$$

for all sufficiently small  $r > 0$ . Thus, as continuity is a local property, we may simplify  $\mathcal{B}$  to

$$\mathcal{B}' := \{B_{\phi_\alpha^* p_\alpha, r}(v) : \alpha \in J, p_\alpha \in \mathcal{P}_\alpha, r > 0, v \in V\},$$

which establishes the local convexity of  $V$  with defining system of seminorms as described within (2.9).  $\square$

Returning to the concrete situation, we thus get the following.

**Corollary 2.25** *Let  $M$  be a manifold and  $k \in \mathbb{N}$ . Then the  $\mathcal{C}^k$ -topology on  $\mathcal{C}^k(M)$  is locally convex with defining system of seminorms given by*

$$p_{K,D}(f) = \max_{p \in K} |Df|, \quad (2.10)$$

*where we vary  $K \subseteq M$  through the compact subsets of  $M$  and  $D \in \text{DiffOp}^k(M)$ .*

Plugging in the local form of differential operators now in principle allows us to return to the chart-based approach we have indicated earlier. Indeed, if  $(U, x)$  is a chart of  $M$  and  $K \subseteq M$  is compact, then localizing  $D_U = D^\alpha \partial_\alpha$  yields the continuous seminorm

$$r_{U,K,D}(f) := \max_{p \in K} \left| D^\alpha \frac{\partial^{|\alpha|} f}{\partial x^\alpha} \right|$$

seminorm on  $\mathcal{C}^k(M)$ . Here, we employ the usual multi-index notation and also Einstein's summation convention. By our considerations, the locally convex topology arising from such seminorms is independent of the chosen atlas and instead intrinsic to the smooth structure.

Finally, the passage to  $\mathcal{C}^\infty(M)$  is based on the simple observation that

$$\mathcal{C}^\infty(M) \hookrightarrow \dots \hookrightarrow \mathcal{C}^{k+1}(M) \hookrightarrow \mathcal{C}^k(M) \hookrightarrow \mathcal{C}^{k-1}(M) \hookrightarrow \dots,$$

i.e. we have canonical linear injections  $\mathcal{C}^\infty(M) \hookrightarrow \mathcal{C}^k(M)$  for all  $k \in \mathbb{N}_0$ . Hence, we may once again use the initial topology to endow  $\mathcal{C}^\infty(M)$  with its own locally convex topology. Taking another look at (2.9), the continuous seminorms are simply given by (2.10), where  $D \in \text{DiffOp}(M)$  is now any differential operator of  $M$ . Note that there are still only *finitely* many derivatives involved in each seminorm. Having established this point of view, the following structural observations are easy to check, but nevertheless instructive.

**Proposition 2.26** *Let  $M$  be a manifold and  $k \in \mathbb{N} \cup \{\infty\}$ .*

- i.) The  $\mathcal{C}^k$ -topology is Fréchet.*
- ii.) The pointwise multiplications*

$$\therefore \mathcal{C}^k \times \mathcal{C}^k \longrightarrow \mathcal{C}^k$$

*are continuous bilinear mappings.*

- iii.) The inclusions  $\mathcal{C}^\ell \hookrightarrow \mathcal{C}^k$  are continuous and dense for all  $0 \leq \ell \leq k$ , where we define*

$$\mathcal{C}^0(M) := \mathcal{C}(M).$$

- iv.) The subspaces*

$$\mathcal{C}_0^\ell(M) := \{f \in \mathcal{C}^\ell(M) : \text{supp } f \text{ is compact}\} \subseteq \mathcal{C}^k(M)$$

*of compactly supported  $\mathcal{C}^\ell$ -functions are dense for all  $\ell = 0, \dots, k$ .*

**Exercise 2.27** Prove Proposition 2.26.

This leaves the question of how to topologize  $\mathcal{C}_0^\ell(M)$  such that it becomes complete itself. If one is interested in distributions (in the sense of generalized functions), this turns out to be an unavoidable question. However, the answer turns out much more involved and ultimately uses final locally convex topologies. Unlike for initial topologies, one then has to demand to stay within the category of locally convex spaces. The interested reader can find the construction of the so-called inductive limit topology within [6, Sec. 4.5–4.6].

### 3 Projective and Injective Tensor Products

There is a plethora of well written treatments of projective tensor products such as [8, §41] and [6, Sec. 15]. The typical discussions of injective tensor products suffer from a distinct lack of focus on seminorms, and as such the lecture notes [17, Sec. 6.3] are the most recommendable source for further reading. We begin our considerations with the central example we would like to understand.

**Example 3.1** We consider the Banach space  $\mathcal{C}([0, 1])$  of complex-valued continuous functions on the unit interval. Its tensor square

$$V := \mathcal{C}([0, 1]) \otimes \mathcal{C}([0, 1])$$

naturally embeds into  $\mathcal{C}([0, 1] \times [0, 1])$  by means of linearly extending

$$(f \otimes g)(x, y) := f(x) \cdot g(y). \quad (3.1)$$

The resulting space constitutes a point separating  $*$ -subalgebra of  $\mathcal{C}([0, 1] \times [0, 1])$  containing the constant functions and as such is dense with respect to the topology of uniform convergence on the unit square  $[0, 1] \times [0, 1]$  by virtue of the Stone-Weierstraß Theorem.

The goal of this section is, roughly speaking, to endow the algebraic tensor product  $V \otimes W$  of two locally convex spaces  $V$  and  $W$  with a locally convex topology derived solely from the topologies of the constituent spaces. Remarkably, one has a multitude of reasonable choices here, a problem famously explored by Grothendieck's [3] during his time as a PhD student. In these notes, we restrict ourselves to injective and projective tensor products, both of which enjoy a plethora of pleasant properties and may be described by explicit defining system of seminorms. Moreover, they pave the way for a conceptually pleasing definition of nuclearity.

We begin with the injective flavour. The principal observation is that for any element  $\varphi \in V'$  of the (continuous) dual space of  $V$ , its absolute value  $|\varphi|$  defines a continuous seminorm on  $V$ . Such seminorms are particularly simple, as they come with a sizable kernel.

**Definition 3.2** *Let  $V$  and  $W$  be locally convex spaces.*

*i.) For  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ , we define their injective tensor product as*

$$(q \otimes_\varepsilon p)(x) := \sup \left\{ |(v' \otimes w')(x)| : v' \in V', |v'| \leq q, w' \in W', |w'| \leq p \right\} \quad (3.2)$$

*for all  $v \in V$  and  $w \in W$ .*

*ii.) The injective tensor product topology on  $V \otimes W$  is generated by the seminorms  $q \otimes_\varepsilon p$ , where we vary  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ . We write  $V \otimes_\varepsilon W$  for the resulting locally convex space.*

The linearity of  $v'$  and  $w'$  ensures that (3.2) indeed constitutes a seminorm. It is straightforward to verify that it suffices to take injective tensor products of seminorms from defining systems of  $V$  and  $W$ . Moreover, the resulting system is indeed filtrating.

**Remark 3.3 (Polars)** Let  $V$  and  $W$  be locally convex spaces. The polar of a subset  $A \subseteq V$  is defined by

$$A^* := \{v' \in V' : |v'(v)| \leq 1 \text{ for all } v \in A\}.$$

The condition on  $v'$  and  $w'$  within (3.2) may thus be rephrased as

$$v' \in B_q^* := (B_{q,1}(0)^{\text{cl}})^* \subseteq V' \quad \text{and} \quad w' \in B_p^* := (B_{p,1}(0)^{\text{cl}})^* \subseteq W',$$

respectively. As the closed cylinders constitute zero neighbourhoods, the Banach-Alaoglu Theorem asserts that these polars are compact with respect to the weak\*-topology, i.e. the topology of pointwise convergence. If  $V$  is normed, then

$$B_{\|\cdot\|}^* = \{v' \in V' : \forall v \in V \ |v'(v)| \leq \|v\|\} = B_1(0)^{\text{cl}},$$

where we endow  $V'$  with the operator norm. This recovers the weak\* compactness of the unit ball in  $V'$ . Thus polars may be regarded as the locally convex incarnation of these balls. More comprehensive discussions can be found within [1].

**Exercise 3.4** Let  $V$  be normed. Interpret the statement of Exercise 2.19 in the context of its bidual  $V''$ .

The following reformulation of (3.2) is often useful, as it only uses one supremum. The idea is that we may pair away a factor of the tensor product to obtain a linear mapping from the continuous dual space to the remaining factors.

**Lemma 3.5** *Let  $V$  and  $W$  be locally convex spaces. Then*

$$\begin{aligned} \iota_V : V \otimes W &\longrightarrow L(V', W), & \iota_V(v \otimes w)v' &:= v'(v) \cdot w \\ \iota_W : V \otimes W &\longrightarrow L(W', V), & \iota_W(v \otimes w)w' &:= w'(w) \cdot v \end{aligned}$$

*extend to well-defined linear injections. Moreover,*

$$(q \otimes_\varepsilon p)(x) = \sup_{|v'| \leq q} p(\iota_V(x)v') = \sup_{|w'| \leq p} q(\iota_W(x)w') \quad (3.3)$$

*for all  $x \in V \otimes W$ ,  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ .*

Note that we have to endow  $V'$  and  $W'$  with locally convex topologies to speak of  $L(V', W)$  and  $L(W', V)$ . As we shall show in a moment, any reasonable choice works, as the mappings are already continuous with respect to the topology of pointwise convergence.

PROOF: The continuity of  $\iota_V(v \otimes w)$  follows from the estimate

$$p(\iota_V(x)v') \leq \sum_k p(\iota_V(v_k \otimes w_k)v') = \sum_k |v'(v_k)| \cdot p(w_k)$$

for all  $v' \in V'$ ,  $p \in \text{cs}(W)$  and  $x = \sum_k v_k \otimes w_k \in V \otimes W$  at once. Indeed, fixing  $x$  and setting

$$r_v : V' \longrightarrow \mathbb{R}, \quad r_v(v') := |v'(v)|,$$

we have shown the continuity estimate

$$p(\iota_V(x)v') \leq \sum_k r_{v_k}(v') \cdot p(w_k)$$

for all  $v' \in V'$  and  $p \in \text{cs}(W)$ . In particular,  $\iota_V(x)$  is continuous for all finer topologies on  $V'$ . The continuity of  $\iota_W(x)$  follows by swapping the roles of  $V$  and  $W$ . Unwrapping (3.3) and using Exercise 2.19, we get

$$\sup_{|v'| \leq q} p(\iota_V(x)v') = \sup_{|v'| \leq q} p\left(\sum_k v'(v_k) \cdot w_k\right)$$

$$\begin{aligned}
&= \sup_{|v'| \leq q} \sup_{|w'| \leq p} \left| w' \left( \sum_k v'(v_k) \cdot w_k \right) \right| \\
&= \sup_{|v'| \leq q} \sup_{|w'| \leq p} \left| \sum_k v'(v_k) \cdot w'(w_k) \right| \\
&= (q \otimes_\varepsilon p)(x).
\end{aligned}$$

The other equality follows analogously.  $\square$

Using this, we may show that  $\otimes_\varepsilon$  formalizes vector-valued versions of many familiar spaces.

**Example 3.6** Let  $M$  be a set and consider the normed space

$$\mathcal{B}(M) := \left\{ f: M \longrightarrow \mathbb{C} : \|f\|_\infty := \sup_{x \in M} |f(x)| < \infty \right\}$$

of all bounded complex-valued mappings defined on  $M$ . Moreover, let  $V$  be locally convex. We are interested in the tensor product  $\mathcal{B}(M) \otimes V$ , which turns out to be related to vector-valued bounded functions on  $M$ . In this context, we call a function  $F: M \longrightarrow V$  bounded if  $F(M) \subseteq V$  is bounded, i.e.  $\sup_{z \in M} q(F(z)) < \infty$  for all  $q \in \text{cs}(V)$ . Hence, the space  $\mathcal{B}(M, V)$  carries a natural locally convex topology induced by the seminorms

$$q_M(F) := \sup_{z \in M} q(F(z)),$$

where we vary  $q \in \text{cs}(V)$ . Notably, we may embed  $\mathcal{B}(M) \otimes V$  into  $\mathcal{B}(M, V)$  by linearly extending

$$(f \otimes v)(z) := f(z) \cdot v. \quad (3.4)$$

The tensor product  $\mathcal{B}(M) \otimes V$  then corresponds precisely of the mappings with images contained in finite dimensional bounded subsets of  $V$ . Using (3.3) and (2.7), we compute

$$\begin{aligned}
(\|\cdot\|_\infty \otimes_\varepsilon q)(x) &= \sup_{|v'| \leq q} \|\iota_V(x)v'\|_\infty \\
&= \sup_{|v'| \leq q} \sup_{z \in M} |\iota_V(x)v'| \\
&= \sup_{z \in M} \sup_{|v'| \leq q} \left| \sum_k v'(v_k) \cdot f_k(z) \right| \\
&= \sup_{z \in M} \sup_{|v'| \leq q} \left| v' \left( \sum_k v_k \cdot f_k(z) \right) \right| \\
&= \sup_{z \in M} q(x(z)) \\
&= q_M(x)
\end{aligned}$$

for  $q \in \text{cs}(V)$  and  $x = \sum_k f_k \otimes v_k \in \mathcal{B}(M) \otimes V$ . Hence, (3.4) extends to a linear embedding

$$\mathcal{B}(M) \otimes_\varepsilon V \hookrightarrow \mathcal{B}(M, V). \quad (3.5)$$

If  $V$  is complete, this embedding has dense range, so that

$$\mathcal{B}(M, V) \cong \mathcal{B}(M) \widehat{\otimes}_\varepsilon V. \quad (3.6)$$

We leave the remaining details as an exercise.

**Exercise 3.7** Let  $M$  be a set and  $V$  a complete Hausdorff locally convex space.

- i.) Prove that  $\mathcal{B}(M, V)$  is complete Hausdorff.
- ii.) Show that (3.5) has dense range.
- iii.) Conclude (3.6).

Instead, we return to Example 3.1.

**Lemma 3.8** *Using the identification (3.1), we have*

$$\mathcal{C}([0, 1]) \widehat{\otimes}_\varepsilon \mathcal{C}([0, 1]) \cong \mathcal{C}([0, 1] \times [0, 1])$$

*as locally convex spaces.*

PROOF: It suffices to prove that the injective tensor product topology translates to the topology of uniform convergence on  $[0, 1] \times [0, 1]$  through the identification (3.1). Let

$$F = \sum_k f_k \otimes g_k \in \mathcal{C}([0, 1]) \otimes \mathcal{C}([0, 1]).$$

As  $\mathcal{C}([0, 1]) \subseteq \mathcal{B}([0, 1])$  is a subspace, we may repeat our computation from Example 3.6 to obtain

$$(\|\cdot\|_\infty \otimes_\varepsilon \|\cdot\|_\infty)(F) = \sup_{x \in [0, 1]} \|F(x, \cdot)\|_\infty = \sup_{x, y \in [0, 1]} |F(x, y)|.$$

Thus (3.1) is an embedding. We have already argued within Example 3.1 that its image is dense by virtue of the Stone-Weierstraß Theorem.  $\square$

**Exercise 3.9** i.) Generalize Lemma 3.8 to  $\mathcal{C}(U)$  for open sets  $U \subseteq \mathbb{R}^n$ .

- ii.) Extend the statement further to smooth functions  $\mathcal{C}^\infty(U)$  for open sets  $U \subseteq \mathbb{R}^n$ .
- iii.) Globalize the assertion to  $\mathcal{C}^\infty(U)$  for open sets  $U \subseteq M$  of a smooth manifold  $M$ .

It is convenient to study the abstract properties of injective and projective tensor products in tandem, so we proceed with yet another definition.

**Definition 3.10 (Projective Tensor Products)** *Let  $V$  and  $W$  be locally convex spaces.*

- i.) *For  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ , we define their projective tensor product as*

$$(q \otimes_\pi p)(x) := \inf \left\{ \sum_k q(v_k) \cdot p(w_k) : x = \sum_k v_k \otimes w_k \right\}. \quad (3.7)$$

- ii.) *The projective tensor product topology on  $V \otimes W$  is generated by the seminorms  $q \otimes_\pi p$ , where we vary  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ . We write  $V \otimes_\pi W$  for the resulting locally convex space.*

As the possible decompositions of a linear combination  $x + \lambda y \in V \otimes W$  are in bijection with the decompositions of  $x$  and  $y$  individually, we get that (3.7) indeed constitute seminorms.

**Exercise 3.11** Let  $V$  and  $W$  be locally convex spaces and  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$ . Prove that (3.2) and (3.7) constitute seminorms on  $V \otimes W$ .

Again, it suffices to use defining systems of seminorms and the resulting system is filtrating. The main reason to study injective and projective tensor products simultaneously is that the latter is always larger than the former.

**Lemma 3.12** ( $\otimes_\pi$  vs.  $\otimes_\varepsilon$ , [16, Cor. of Prop. 43.4]) *Let  $V$  and  $W$  be locally convex spaces. Then the identity mapping*

$$V \otimes_\pi W \longrightarrow V \otimes_\varepsilon W$$

*is continuous. More precisely, if  $p \in \text{cs}(V)$  and  $q \in \text{cs}(W)$ , then*

$$p \otimes_\varepsilon q \leq p \otimes_\pi q. \quad (3.8)$$

PROOF: Let  $x \in V \otimes W$  and consider a finite decomposition

$$x = \sum_k v_k \otimes w_k$$

into factorizing tensors as well as  $v' \in B_p^*$  and  $w' \in B_q^*$ . Then

$$|(v' \otimes w')(x)| \leq \sum_k |v'(v_k) \cdot w'(w_k)| \leq \sum_k p(v_k) \cdot q(w_k).$$

Note that the left-hand side is independent of the chosen decomposition, whereas the right-hand side does not depend on the choice of  $v'$  and  $w'$ . Thus, taking the infimum over all decompositions of  $x$  and the supremum over all  $v'$  and  $w'$  in the respective polars proves the continuity estimate (3.8).  $\square$

Due to the supremum within (3.2) and infimum in (3.7), it is typically rather cumbersome to compute projective and injective tensor products explicitly. The triangle inequality combined with the following Lemma provides a simple yet powerful estimate from above.

**Lemma 3.13** *Let  $V$  and  $W$  be locally convex spaces. Then*

$$(p \otimes_\varepsilon q)(v \otimes w) = p(v) \cdot q(w) = (p \otimes_\pi q)(v \otimes w) \quad (3.9)$$

*for all  $p \in \text{cs}(V)$ ,  $q \in \text{cs}(W)$ ,  $v \in V$  and  $w \in W$ .*

That is to say, projective and injective tensor products factorize on factorizing tensors. The proof is based on the Hahn-Banach Theorem and showcases the interplay between both concepts.

PROOF (OF LEMMA 3.13): Let  $p \in \text{cs}(V)$ ,  $q \in \text{cs}(W)$ ,  $v \in V$  and  $w \in W$ . By (3.7), we have

$$(p \otimes_\pi q)(v \otimes w) \leq p(v) \cdot q(w)$$

as  $v \otimes w$  already comes with a decomposition into factorizing tensors out of the box. For the converse inequality, we consider the linear functionals

$$v': \text{span } v \longrightarrow \mathbb{C}, \quad v'(\lambda \cdot v) := \lambda \cdot p(v)$$

$$w': \text{span } w \longrightarrow \mathbb{C}, \quad w'(\lambda \cdot w) := \lambda \cdot q(w),$$

which fulfil  $|v'| \leq p$  and  $|w'| \leq q$  by construction. Invoking the Hahn-Banach Theorem ??, we may thus extend both continuously to all of  $V$  and  $W'$  while preserving the continuity estimates. That is to say,  $v' \in V'$  and  $w' \in W'$  get to partake in the supremum within (3.2), which yields the estimate

$$(p \otimes_\varepsilon q)(v \otimes w) \geq \left| (v' \otimes w')(v \otimes w) \right| = |v'(v) \cdot w'(w)| = p(v) \cdot q(w).$$

Putting everything together and using (3.8), we arrive at

$$p(v) \cdot q(w) \geq (p \otimes_\pi q)(v \otimes w) \geq (p \otimes_\varepsilon q)(v \otimes w) \geq p(v) \cdot q(w),$$

which is (3.9). □

Viewing (3.9) as a continuity estimate, the following is immediate.

**Corollary 3.14** *Let  $V$  and  $W$  be locally convex spaces. Then*

$$\otimes_\varepsilon: V \times W \longrightarrow V \otimes_\varepsilon W \quad \text{and} \quad \otimes_\pi: V \times W \longrightarrow V \otimes_\pi W$$

*are continuous bilinear mappings.*

Another natural question is whether tensor products of norms are again norms. In view of the infimum within (3.7), this is not completely obvious if done directly. That being said, (3.8) ensures that it suffices to provide a lower bound for the injective tensor product (3.2). Here, the supremum in the definition comes to our aid much like in the proof of Lemma 3.13. This is a common theme. More generally, this question may be recast as the inheritance of the Hausdorff property upon taking tensor products. The precise statement is the following.

**Proposition 3.15 (Hausdorff Property)** *Let  $V$  and  $W$  be locally convex spaces.*

- i.) If  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$  are norms, then so are  $q \otimes_\varepsilon p$  and  $q \otimes_\pi p$ .*
- ii.) The tensor products  $V \otimes_\varepsilon W$  and  $V \otimes_\pi W$  are Hausdorff if  $V$  and  $W$  are. The converse holds whenever  $V \neq \{0\} \neq W$ .*

PROOF: We begin with the second statement and assume first that  $V$  and  $W$  are Hausdorff. Let  $x = \sum_{k=1}^N v_k \otimes w_k \in V \otimes W$ . Without loss of generality, we may assume

$$v_1 \notin \text{span}\{v_2, \dots, v_N\};$$

otherwise we write  $v_1$  as a linear combination of the remaining vectors and simplify back to multiples of  $v_2, \dots, v_N$ . Relabeling and repeating the process then terminates after finitely many steps, achieving the desired condition. By assumption, we find  $q \in \text{cs}(V)$  with  $q(v_1) > 0$  and  $p \in \text{cs}(W)$  with  $p(w_1) > 0$ . Similar to before, we consider the linear functionals

$$v': \text{span}\{v_1, \dots, v_N\} \longrightarrow \mathbb{C}, \quad v'\left(\sum_{k=1}^N \lambda_k v_k\right) := \lambda \cdot q(v_1)$$

$$w': \text{span } w_1 \longrightarrow \mathbb{C}, \quad w'(\lambda \cdot w_1) := \lambda \cdot p(w_1).$$

Again,  $|v'| \leq q$  and  $|w'| \leq p$ , and we use the Hahn-Banach Theorem ?? to continuously extend both to  $V$  and  $W$ , respectively, while preserving the continuity estimates. Together with (3.8), we arrive at

$$(q \otimes_\pi p)(x) \geq (q \otimes_\varepsilon p)(x) \geq |(v' \otimes w')(x)| = \left| \sum_{k=1}^N \underbrace{v'(v_k)}_{=\delta_{k,1} \cdot q(v_1)} w'(w_k) \right| = q(v_1) \cdot p(w_1) > 0.$$

Varying  $x$  establishes the Hausdorff property of  $V \otimes_\pi W$  and  $V \otimes_\varepsilon W$ . Assume conversely that neither  $V$ , nor  $W$  are the zero space and  $V \otimes_\pi W$  or  $V \otimes_\varepsilon W$  is Hausdorff. By choosing some  $v \in V \setminus \{0\}$ , we get a linear injection

$$\iota_v: W \longrightarrow V \otimes W, \quad \iota_v(w) := v \otimes w.$$

By virtue of Lemma 3.13, it even constitutes a topological embedding for either topology. That is to say, it is a homeomorphism onto its image. Hence,  $W$  inherits the Hausdorff property from the ambient space. The argument for  $V$  is analogous. This completes the proof of *ii.*) and we turn towards *i.*). To this end, assume that  $q \in \text{cs}(V)$  and  $p \in \text{cs}(W)$  are norms. Then we may regard  $(V, q)$  and  $(W, p)$  as normed spaces, whose locally convex topology is generated by

$$\{r \cdot q: r > 0\} \quad \text{resp.} \quad \{r \cdot p: r > 0\}.$$

As tensor products of Hausdorff spaces are Hausdorff by *ii.*), we get that the defining system

$$\{(r_1 \cdot q) \otimes_{\pi/\varepsilon} (r_2 \cdot p): r_1, r_2 > 0\} = \{r \cdot (q \otimes_{\pi/\varepsilon} p): r > 0\}$$

has no joint kernel. But this means

$$q \otimes_{\pi/\varepsilon} p$$

is a norm itself, completing the proof.  $\square$

It is instructive to make the argument we have just used precise.

**Exercise 3.16** Let  $(V, \|\cdot\|)$  be a normed space. Prove that the system of seminorms

$$\{r \cdot \|\cdot\|: r > 0\}$$

is filtrating and that the associated locally convex topology reproduces the norm-topology.

Another pleasant property of both flavours of tensor products is that naively gluing continuous linear mappings preserves continuity. We begin with the injective version.

**Lemma 3.17** *Let  $\phi_j: V_j \longrightarrow W_j$  be continuous linear mappings between locally convex spaces for  $j = 1, 2$ . Then the mapping*

$$\phi_1 \otimes \phi_2: V_1 \otimes_\varepsilon V_2 \longrightarrow W_1 \otimes_\varepsilon W_2$$

*defined by linear extension of*

$$(\phi_1 \otimes \phi_2)(v_1 \otimes v_2) := \phi_1(v_1) \otimes \phi_2(v_2)$$

*is continuous.*

PROOF: Let  $p_1 \in \text{cs}(W_1)$  and  $p_2 \in \text{cs}(W_2)$ . By continuity of  $\phi_1$  and  $\phi_2$ , there exist corresponding seminorms  $q_1 \in \text{cs}(V_1)$  and  $q_2 \in \text{cs}(V_2)$  such that

$$\phi_1^* p_1 = p_1 \circ \phi_1 \leq q_1 \quad \text{and} \quad \phi_2^* p_2 = p_2 \circ \phi_2 \leq q_2.$$

Consequently, we have

$$B_{\phi^* p_1}^* \subseteq B_{q_1}^* \quad \text{and} \quad B_{\phi^* p_2}^* \subseteq B_{q_2}^*.$$

Writing  $\phi := \phi_1 \otimes \phi_2$ , this implies

$$\begin{aligned} (p_1 \otimes_\varepsilon p_2)(\phi(x)) &= \sup_{w'_1 \in B_{p_1}^*} \sup_{w'_2 \in B_{p_2}^*} \left| (w'_1 \otimes w'_2)(\phi(x)) \right| \\ &= \sup_{w'_1 \in B_{p_1}^*} \sup_{w'_2 \in B_{p_2}^*} \left| (\phi_1^* w'_1 \otimes \phi_2^* w'_2)(x) \right| \\ &\leq \sup_{v'_1 \in B_{q_1}^*} \sup_{v'_2 \in B_{q_2}^*} \left| (v'_1 \otimes v'_2)(x) \right| \\ &\leq (q_1 \otimes_\varepsilon q_2)(x), \end{aligned}$$

which completes the proof.  $\square$

**Exercise 3.18** Prove that taking injective and projective tensor products is associative up to the usual linear algebraic identifications. That is to say, show that

$$p_1 \otimes_{\pi/\varepsilon} (p_2 \otimes_{\pi/\varepsilon} p_3) = (p_1 \otimes_{\pi/\varepsilon} p_2) \otimes_{\pi/\varepsilon} p_3$$

as mappings on the triple tensor product  $V_1 \otimes V_2 \otimes V_3$  of locally convex spaces  $V_1, V_2, V_3$  and  $p_j \in \text{cs}(V_j)$  for  $j = 1, 2, 3$ . Conclude that the resulting locally convex spaces are canonically linearly homeomorphic.

Recall the universal property of the algebraic tensor product: Given a bilinear mapping

$$\phi: V \times W \longrightarrow X,$$

there exists a unique linear mapping

$$\Phi: V \otimes W \longrightarrow X \quad \text{such that } \Phi \circ \otimes = \phi.$$

Our Corollary 3.14 ensures that the continuity of  $\Phi$  implies the continuity of  $\phi$ . The natural question is thus whether the converse also holds. This turns out to be a distinguishing property of the projective tensor product, which we could have used as the definition instead. This statement is known as the *infimum argument* and an ubiquitous tool within strict deformation quantization as discussed in the survey [18]. Using the associativity from Exercise 3.18, the general statement is the following.

**Proposition 3.19 (Infimum argument, [16, Prop. 43.4])** *Let  $V_1, \dots, V_n, W$  be locally convex spaces and*

$$\phi: V_1 \times \dots \times V_n \longrightarrow W$$

*be  $n$ -linear with corresponding linear map*

$$\Phi: V_1 \otimes \dots \otimes V_n \longrightarrow W.$$

Endow  $V_1 \times \cdots \times V_n$  with the Cartesian product topology and  $V_1 \otimes \cdots \otimes V_n$  with the projective tensor product topology. Then  $\phi$  is continuous if and only if  $\Phi$  is. More precisely, if for a continuous seminorm  $q \in \text{cs}(W)$  there are  $p_1 \in \text{cs}(V_1), \dots, p_n \in \text{cs}(V_n)$  such that

$$q(\phi(v_1, \dots, v_n)) \leq p_1(v_1) \cdots p_n(v_n) \quad \text{for all } v_1 \in V_1, \dots, v_n \in V_n, \quad (3.10)$$

then

$$q(\Phi(v)) \leq (p_1 \otimes \cdots \otimes p_n)(v) \quad \text{for all } v \in V_1 \otimes \cdots \otimes V_n, \quad (3.11)$$

and vice versa.

PROOF: It suffices to treat the case  $n = 2$ . As already noted, we may view Lemma 3.13 as the continuity estimate

$$(p_1 \otimes_\pi p_2)(v_1 \otimes v_2) \leq p_1(v_1) \cdot p_2(v_2).$$

Hence, (3.10) implies (3.11) by virtue of  $\Phi \circ \otimes_\pi = \phi$ . Assume conversely (3.11) and let

$$x = \sum_k v_k \otimes w_k \in V_1 \otimes V_2.$$

Then we have

$$q(\Phi(x)) \leq \sum_k q(\Phi(v_k \otimes w_k)) = \sum_k q(\phi(v_k, w_k)) \leq \sum_k p_1(v_k) \cdot p_2(w_k).$$

Taking the infimum over all decompositions of  $x$  into factorizing tensors thus yields

$$q(\Phi(x)) \leq (p_1 \otimes_\pi p_2)(x).$$

Varying  $x \in V_1 \otimes V_2$  thus establishes (3.11).  $\square$

The upshot is that it suffices to prove continuity estimates for multilinear mappings on factorizing tensors. As a simple application, we may prove the analogue of Lemma 3.17 for projective tensor products.

**Corollary 3.20** *Let  $\phi_j: V_j \longrightarrow W_j$  be continuous linear mappings between locally convex spaces for  $j = 1, 2$ . Then the mapping*

$$\phi_1 \otimes \phi_2: V_1 \otimes_\pi V_2 \longrightarrow W_1 \otimes_\pi W_2$$

*defined by linear extension of*

$$(\phi_1 \otimes \phi_2)(v_1 \otimes v_2) := \phi_1(v_1) \otimes \phi_2(v_2)$$

*is continuous.*

PROOF: By Proposition 3.19 and Corollary 3.14, it suffices to prove the continuity of the bilinear mapping

$$\phi: V_1 \times V_2 \longrightarrow W_1 \times W_2, \quad \phi(v_1, v_2) := (\phi_1(v_1), \phi_2(v_2)).$$

By continuity of  $\phi_1$  and  $\phi_2$  and the definition of the product topology, this is obvious.  $\square$

**Example 3.21** Let  $V$  be a complete Hausdorff locally convex space. A sequence of vectors  $\gamma := (v_n) \subseteq V$  is called absolutely summable if

$$q(\gamma) := \sum_{n=0}^{\infty} q(v_n) < \infty \quad (3.12)$$

for all  $q \in \text{cs}(V)$ . We write  $\ell^1[V]$  for the corresponding locally convex space. The usual  $\ell^1$ -space then arises for  $V = (\mathbb{C}, |\cdot|)$ , viewed as a locally convex space. We claim that

$$\ell^1 \widehat{\otimes}_{\pi} V \cong \ell^1[V] \quad (3.13)$$

as locally convex spaces. To see this, we consider the mapping

$$\phi: \ell^1 \widehat{\otimes}_{\pi} V \longrightarrow \ell^1[V]$$

defined by linear extension of

$$(\phi(a \otimes v))_n := a_n \cdot v.$$

By virtue of

$$q(\phi(a \otimes v)) = \sum_{n=0}^{\infty} q(a_n \cdot v) = \sum_{n=0}^{\infty} |a_n| \cdot q(v) = \|a\|_1 \cdot q(v) \quad (3.14)$$

for all  $a \in \ell^1$  and  $v \in V$  and Proposition 3.19, this is indeed well-defined on all of the completion. Moreover, the resulting mapping is clearly injective on  $\ell^1 \otimes_{\pi} V$  and thus the same is true for its unique continuous linear extension. Let now  $\gamma \in \ell^1[V]$ . We check that the series

$$\Gamma := \sum_{n=0}^{\infty} e_n \otimes \gamma_n$$

with the usual unit sequences  $e_n \in \ell^1$  given by  $e_n(k) := \delta_{n,k}$  converges within  $\ell^1 \widehat{\otimes}_{\pi} V$ . Indeed, by Lemma 3.13 we have

$$\sum_{n=0}^{\infty} (\|\cdot\|_1 \otimes_{\pi} q)(e_n \otimes \gamma_n) = \sum_{n=0}^{\infty} \|e_n\|_1 \cdot q(\gamma_n) = \sum_{n=0}^{\infty} q(\gamma_n) = q(\gamma) < \infty$$

for all  $q \in \text{cs}(V)$ . By completeness, we know that absolute convergence implies convergence, and thus  $\Gamma \in \ell^1 \widehat{\otimes}_{\pi} V$  is well-defined. By continuity of  $\phi$  and the completeness of  $V$ , this implies the convergence of

$$\phi(\Gamma) = \sum_{n=0}^{\infty} \phi(e_n \otimes \gamma_n) = \sum_{n=0}^{\infty} e_n \cdot \gamma_n = \gamma.$$

Hence,  $\phi$  is also surjective and thus bijective. Finally, reading (3.14) backwards asserts the continuity of the inverse mapping. Note that passing to the completion was essential. The image

$$\phi(\ell^1 \otimes_{\pi} V)$$

only consists of sequences living within finite dimensional subspaces of  $V$ . As a particular consequence of the equality (3.14) combined with Proposition 3.19, we get that

$$\ell^1 \widehat{\otimes}_\pi \ell^1[J] \cong \ell^1[\mathbb{N}_0 \times J]$$

for any index set  $J$ . More generally, one may use the same methods to prove

$$\ell^1[I] \widehat{\otimes}_\pi \ell^1[J] \cong \ell^1[I \times J].$$

Finally, we note that there exist measure theoretic generalizations, which can be found within [6, Sec. 16.7]. It should be noted that this particularly nice compatibility is particular to  $\ell^1$  and fails e.g. for the space of zero sequences, see [5, Ex. 2.6.2] for further discussion. Another such example is the Hilbert space  $\ell^2$ , where one should use the Hilbert tensor product instead.

## 4 Nuclearity

The de-facto source regarding nuclearity remains [12], even though some of the treatment has become dated. A more modern point of view can be found in the lecture notes [17].

In Lemma 3.12, we have seen that there is a canonical continuous linear mapping

$$V \otimes_\pi W \longrightarrow V \otimes_\varepsilon W$$

between the projective and injective tensor products of two locally convex spaces  $V$  and  $W$ . It turns out that requiring this mapping to even constitute a homeomorphism has far reaching consequences. This is the notion of nuclearity.

**Definition 4.1 (Nuclearity)** *We call a locally convex space  $V$  nuclear if the canonical mapping*

$$V \otimes_\pi W \longrightarrow V \otimes_\varepsilon W$$

*constitutes a homeomorphism for all locally convex spaces  $W$ .*

In the sequel, we may thus suppress the index for the tensor product, whenever we work with nuclear spaces.

From the Definition, it is clear that we need some better criteria to check nuclearity.

**Proposition 4.2 ...**

There are many more equivalent, yet different looking and thus situationally useful characterizations of nuclearity. Our next goal is to generalize the Riemann rearrangement Theorem to nuclear spaces. We have already met the space of absolutely summable sequences  $\ell^1[V]$  of a complete Hausdorff space within Example 3.21. We managed to identify it as the completion of the projective tensor product of  $\ell^1$  and  $V$  within (3.13). We prove the analogous result holds for injective tensor products and unconditionally convergent series.

**Example 4.3 (Unconditional Summability I)** Let  $V$  be a complete Hausdorff locally convex space and  $(v_n)_n \subseteq V$  a sequence. Recall that the corresponding series

$$\sum_{n=0}^{\infty} v_n$$

is called unconditionally convergent to  $v \in V$  if

$$\sum_{n=0}^{\infty} v_{\sigma(n)} = v$$

for all bijections  $\sigma: \mathbb{N}_0 \rightarrow \mathbb{N}_0$ . That is to say, one may rearrange the series without spoiling convergence or altering the limit. We write  $\ell^1(V)$  for the resulting vector space. At this point, it is not completely clear how one should topologize  $\ell^1(V)$ . Note first that the canonical mapping

$$\phi: \ell^1 \otimes V \rightarrow \ell^1(V), \quad (\phi(\gamma \otimes v))_n := \gamma_n \cdot v \quad (4.1)$$

constitutes a linear injection. Indeed, that this sequence is indeed unconditionally summable by the usual Riemann Rearrangement Theorem, as every tensor is a finite linear combination of factorizing tensors, and thus the resulting limit is taken within a finite dimensional subspace of  $V$ . Invoking once again (3.3), we compute

$$\begin{aligned} (\|\cdot\|_1 \otimes_{\varepsilon} q)(x) &= \sup_{|v'| \leq q} \|\iota_V(x)v'\|_1 \\ &= \sup_{|v'| \leq q} \left\| \sum_k v'(v_k) \cdot \gamma_k \right\|_1 \\ &= \sup_{|v'| \leq q} \sum_{n=0}^{\infty} \left| \sum_k v'(v_k) \cdot \gamma_k(n) \right| \\ &= \sup_{|v'| \leq q} \sum_{n=0}^{\infty} \left| v' \left( \sum_k v \cdot \gamma_k(n) \right) \right| \\ &= \sup_{|v'| \leq q} \sum_{n=0}^{\infty} |v'(x_n)| \end{aligned}$$

for  $q \in \text{cs}(V)$  and  $x = \sum_k \gamma_k \otimes v_k \in \ell^1 \otimes V$ , where we write

$$x_n := \sum_k (\gamma_k)(n) \cdot v_k \in V \quad \text{for all } n \in \mathbb{N}_0.$$

Hence, a reasonable choice of seminorms turns out to be

$$q_w((v_n)_n) := \sup_{|v'| \leq q} \sum_{n=0}^{\infty} |v'(v_n)|, \quad (4.2)$$

where we vary  $q \in \text{cs}(V)$ . The index  $w$  alludes to weak convergence. It is not completely obvious that the suprema within (4.2) are finite. To see this, we take a detour to note the following characterization of unconditional convergence.

**Lemma 4.4** *Let  $V$  be a Hausdorff locally convex space and  $(v_n)_n \subseteq V$  a sequence. Then the series  $\sum_{n \in \mathbb{N}_0} v_n$  converges unconditionally to  $v \in V$  iff for every  $\varepsilon > 0$  and  $q \in \text{cs}(V)$  there exists a finite set  $F_0 \subseteq \mathbb{N}_0$  such that*

$$q\left(v - \sum_{n \in F} v_n\right) \leq \varepsilon \quad (4.3)$$

for all finite sets  $F_0 \subseteq F \subseteq \mathbb{N}_0$ .

PROOF: Assume first that the condition from the Lemma holds and let  $\sigma: \mathbb{N}_0 \rightarrow \mathbb{N}_0$  be a bijection. Moreover, let  $\varepsilon > 0$  and  $q \in \text{cs}(V)$ . By assumption, we find some finite set  $F_0 \subseteq \mathbb{N}_0$  such that (4.3) holds for all finite sets  $F_0 \subseteq F \subseteq \mathbb{N}_0$ . As  $F_0$  is finite, there exists some  $N_0 \in \mathbb{N}_0$  such that  $\sigma(n) \geq \max F_0$  for all  $n \geq N_0$ . That is to say by bijectivity of  $\sigma$ , we have

$$F_0 \subseteq \sigma(\{1, \dots, N_0\}),$$

which implies

$$q\left(v - \sum_{n=0}^N v_{\sigma(n)}\right) \leq \varepsilon$$

for all  $N \geq N_0$ . Varying  $\varepsilon$  and  $q$  establishes the convergence of  $\sum_{n=0}^{\infty} v_{\sigma(n)}$  towards  $v$ .

Assume, conversely, that the condition stated in the Lemma fails. Then there exists some constant  $\varepsilon > 0$  and  $q \in \text{cs}(V)$  such that for every finite set  $F_0 \subseteq \mathbb{N}_0$ , there exists another finite set  $F_0 \subseteq F \subseteq \mathbb{N}_0$  such that

$$q\left(v - \sum_{n \in F} v_n\right) > \varepsilon.$$

We use this to define a bijection  $\sigma: \mathbb{N}_0 \rightarrow \mathbb{N}_0$  as follows: first, we set  $\sigma(0) := 0$ . If now  $\sigma(0), \dots, \sigma(n)$  are already defined for some  $n \in \mathbb{N}_0$ , then take  $F_0 := \{\sigma(0), \dots, \sigma(n)\}$  with corresponding finite set  $F_0 \subseteq F \subseteq \mathbb{N}_0$ . If  $F = F_0$ , then we set

$$\sigma(n+1) := \min(\mathbb{N}_0 \setminus F_0).$$

Otherwise, we define

$$\sigma(n+1), \sigma(n+2), \dots, \sigma(n + |F \setminus F_0|)$$

as the elements of  $F \setminus F_0$  in ascending order. Note that the finite sets  $F$  within each step contain all the preceding ones. Inductively, this defines a bijection  $\sigma$ . We check that the accordingly rearranged series does not converge to  $v$ . To this end, let  $N_0 \in \mathbb{N}_0$ . Then, by construction of  $\sigma$ , there exists some  $N_1 \geq N_0$  such that

$$F(N_0) = \{\sigma(0), \dots, \sigma(N_1)\}.$$

This implies the desired inequality

$$q\left(v - \sum_{n=0}^{N_1} v_{\sigma(n)}\right) > \varepsilon.$$

**Example 4.5 (Unconditional Summability II)** Let  $V$  be a complete Hausdorff locally convex space. We return to the finiteness of (4.2) for unconditionally summable sequences  $(v_n)_n \subseteq V$  with limit  $v$ . Fix  $v' \in V$ . Invoking Lemma 4.4, we find some finite set  $F_0 \subseteq \mathbb{N}_0$  such that

$$\left| v'(v) - \sum_{n \in F} v'(v_n) \right| = \left| v' \left( v - \sum_{n \in F} v_n \right) \right| \leq 1$$

for all finite sets  $F_0 \subseteq F \subseteq \mathbb{N}_0$ , where we use that  $|v'| \in \text{cs}(V)$  by continuity of  $v'$ . Consequently,

$$\sum_{n \in J} |v'(v_n)| \leq 1 + |v'(v)|$$

for all finite sets  $F_0 \subseteq F \subseteq \mathbb{N}_0$ . Hence, the series

$$\sum_{n=0}^{\infty} v'(v_n)$$

converges absolutely within  $\mathbb{C}$ , as the corresponding sequence of partial sums is bounded from above and monotonically increasing. Hence, each individual term within the supremum (4.2) is finite. The remainder of the proof uses technology beyond the scope of these notes. For the sake of completeness, we give the argument regardless. Consider the sets

$$\Phi_r := \left\{ v' \in V' : \sum_{n=0}^{\infty} |v'(v_n)| \leq r \right\} = \bigcap_{J \in \mathcal{F}} \left\{ v' \in V' : \sum_{n \in J} |v'(v_n)| \leq r \right\} \quad (4.4)$$

for all  $r > 0$  and where  $\mathcal{F}$  denotes the collection of all finite subsets of  $\mathbb{N}_0$ . By what we have just shown, we know

$$\bigcup_{r>0} \Phi_r = \bigcup_{r>0} r \cdot \Phi_1 = V'.$$

Moreover,  $\Phi_r$  is absolutely convex and closed with respect to the topology of pointwise convergence on  $V'$ . Indeed, the closedness of each of the members of the intersection in (4.4) follows from the finiteness of  $J$  and thus  $\Phi_r$  is closed as an intersection of closed sets. We have shown that each  $\Phi_r$ , in particular  $\Phi_1$ , constitutes a *barrel* within  $V'$ . Hence, pulling back the intersection  $\Phi_1 \cap B_q^*$  to functionals on the local Banach space

$$V_q := \widehat{V / \ker q}$$

with the obvious quotient norm  $[q]$  yields a barrel  $[\Phi] \subseteq V'_q$ . As a Banach space,  $V_q$  is barrelled and thus  $[\Phi]$  constitutes a zero neighbourhood and is thus contained in some sufficiently large norm-ball. That is to say, there exists  $r > 0$  such that

$$\sum_{n=0}^{\infty} |v'(v_n)| = \sum_{n=0}^{\infty} |[v]'([v_n])| \leq r \quad \text{for all } v' \in B_q^*,$$

where we have used that  $\ker q \subseteq \ker v'$ . Hence, (4.2) is indeed well-defined. In passing, we note that our argument did not rely on the unconditional convergence and may be readily generalized to arbitrary summable sequences. This gives rise to yet another complete space of sequences on  $V$ , which constitutes yet another contender for which space

the symbol  $\ell^1(V)$  should really denote. Within these notes, it shall unnamed and thus shrouded in mystery. Our next goal is to show

$$\ell^1 \widehat{\otimes}_\varepsilon V \cong \ell^1(V), \quad (4.5)$$

where we once again use the assumed completeness of  $V$ . As in the projective situation, we establish the density of the inclusion (4.1). Let  $(v_n)_n \in \ell^1(V)$  with  $v := \sum_{n=0}^\infty v_n$  and consider the sequence  $(\gamma_m)$  given by

$$\gamma_m := \sum_{n=0}^m e_n \otimes v_n \in \ell^1 \otimes_\varepsilon V.$$

Given  $\varepsilon > 0$  and  $q$ , we invoke Lemma 4.4 to find a finite set  $F_0 \subseteq \mathbb{N}_0$  such that

$$q\left(v - \sum_{n \in F} v_n\right) \leq \varepsilon$$

for all finite sets  $F_0 \subseteq F \subseteq \mathbb{N}_0$ . Obversely, this means that

$$q\left(\sum_{n \in F} v_n\right) \leq q\left(v - \sum_{n \in F_0} v_n\right) + q\left(v - \sum_{n \in F \cup F_0} v_n\right) \leq 2\varepsilon$$

for any finite set  $F \subseteq \mathbb{N}_0$  with  $F \cap F_0 = \emptyset$ . In particular, we get

$$\left|v'\left(\sum_{n \in F} v_n\right)\right| = \left|\sum_{n \in F} v'(v_n)\right| \leq 2\varepsilon$$

for any finite set  $F \subseteq \mathbb{N}_0$  with  $F \cap F_0 = \emptyset$  and  $v' \in V'$  with  $|v'| \leq q$ . Covering  $\mathbb{N}_0$  with the sets

$$\mathbb{N}_\pm^\pm := \{n \in \mathbb{N}_0 : \pm \operatorname{Re} v'(v_n) \geq 0, \pm \operatorname{Im} v'(v_n) \geq 0\}$$

leads to the estimate

$$\begin{aligned} \sum_{n \in F} |v'(v_n)| &\leq \sum_{n \in F} |\operatorname{Re} v'(v_n)| + \sum_{n \in F} |\operatorname{Im} v'(v_n)| \\ &= \sum_{n \in F \cap \mathbb{N}_+^+} |\operatorname{Re} v'(v_n)| + \sum_{n \in F \cap \mathbb{N}_-^+} |\operatorname{Re} v'(v_n)| \\ &\quad + \sum_{n \in F \cap \mathbb{N}_+^-} |\operatorname{Im} v'(v_n)| + \sum_{n \in F \cap \mathbb{N}_-^-} |\operatorname{Im} v'(v_n)| \\ &\leq 8\varepsilon \end{aligned}$$

for the same data and where we have used the triangle inequality in the plane to infer

$$|z| = |\operatorname{Re} z + \operatorname{Im} z| \leq |\operatorname{Re} z| + |\operatorname{Im} z| \quad \text{for all } z \in \mathbb{C}.$$

Varying the data, we get

$$\sup_{|v'| \leq q} \sum_{n > N} |v'(v_n)| \leq 8\varepsilon.$$

By construction, we have on the other hand

$$v_n - \gamma_m(n) = \begin{cases} 0 & \text{for } n = 0, \dots, m, \\ v_n & \text{for } n > m, \end{cases}$$

and thus

$$q_w((v_n)_n - \gamma_m) = \sup_{|v'| \leq q} \sum_{n=0}^{\infty} |v'(v_n - \gamma_m(n))| = \sup_{|v'| \leq q} \sum_{n=m+1}^{\infty} |v'(v_n)| \leq 8\varepsilon$$

for all  $m \geq N$ . This proves the desired convergence  $\gamma_m \rightarrow (v_n)_n$ .

Once again, it remains to establish the completeness of the ambient space to complete the argument.

**Exercise 4.6** Let  $V$  be a complete Hausdorff locally convex space. Prove that the space  $\ell^1(V)$  of unconditionally summable sequences is complete Hausdorff.

In passing, we note the elementary estimate

$$q_w((v_n)_n) = \sup_{|v'| \leq q} \sum_{n=0}^{\infty} |v'(v_n)| \leq \sum_{n=0}^{\infty} q(v_n) = q((v_n)_n) \quad \text{for all } (v_n)_n \in \ell^1[V],$$

with the seminorms for  $\ell^1[V]$  from (3.12). In view of (3.13) and (4.5) this is nothing but a particular case of (3.8). We may encode this as a canonical linear injection

$$\ell^1[V] \longrightarrow \ell^1(V).$$

Putting everything together, we arrive at the following Theorem.

**Theorem 4.7 (Summability in Nuclear spaces)** *Let  $V$  be a complete Hausdorff nuclear locally convex space. Then  $\ell^1[V] \cong \ell^1(V)$ . That is to say, any series within  $V$  converges absolutely iff it converges unconditionally.*

PROOF: Combining the very definition of nuclearity with (3.13) and (4.5) yields

$$\ell^1[V] \cong \ell^1 \widehat{\otimes}_{\pi} V \cong \ell^1 \widehat{\otimes}_{\varepsilon} V \cong \ell^1(V).$$

The additional statement follows from unwrapping the canonical homeomorphisms.  $\square$

This constitutes a direct generalization of Riemann's Rearrangement Theorem beyond the finite-dimensional situation. Note that the completeness is immaterial and may always be remedied by passing to the completion of  $V$  as a preliminary step. Remarkably, the equality of unconditional and absolute summability characterizes nuclearity by [?]. However, this means that infinite-dimensional normed spaces are never nuclear.

**Example 4.8** Consider  $V := \ell^2$  and write  $e_n$  for the usual orthogonal basis with entries given by  $e_n(k) = \delta_{n,k}$  for all  $n, k \in \mathbb{N}$ . Then the series

$$\sum_{n=1}^{\infty} \frac{e_n}{n}$$

converges unconditionally, but not absolutely. Hence,  $\ell^2$  is not nuclear by Theorem 4.7. By virtue of the Riesz-Fischer Theorem, this extends to any infinite dimensional Hilbert space. In fact, the same is true for any infinite dimensional *normed* space by an application of the highly non-trivial Dvoretzky-Rogers Theorem [2].

We proceed with pursuing another remarkable property of nuclear spaces: the existence of compact sets with non-empty open interior. That is to say, we are going to prove a variant of Montel's Theorem for nuclear spaces. In passing, we introduce quasicompleteness, which is a weaker notion of completeness useful in many applications. Recall that every Cauchy sequence is bounded. The same need not be true about Cauchy nets. It is instructive to play with this within some examples.

**Exercise 4.9** Construct a unbounded Cauchy net. Argue that your net converges within the completion of your space, yet remains unbounded there.

This leads to the insight that unbounded Cauchy nets might not be the ones we care about. Accordingly, there is a more appropriate flavour of completeness.

**Definition 4.10 (Quasicompleteness)** *A locally convex space  $V$  is called quasicomplete if every bounded Cauchy net converges.*

**Theorem 4.11** *Let  $V$  be a nuclear as well as quasicomplete space. Then every bounded subset  $B \subseteq V$  has compact closure.*

PROOF: This is an elaborate exercise for another day. □

## References

- [1] CONWAY, J. B.: *A Course in Functional Analysis*, vol. 96 in *Graduate Texts in Mathematics*. Springer-Verlag, Heidelberg, Berlin, New York, 2. edition, 1990. 1
- [2] DVORETZKY, A., ROGERS, C. A.: *Absolute and unconditional convergence in normed linear spaces*. Proc. Nat. Acad. Sci. U.S.A. **36** (1950), 192–197. 34
- [3] GROTHENDIECK, A.: *Produits tensoriels topologiques et espaces nucléaires*. Mem. Amer. Math. Soc. **16** (1955), Chapter 1: 196 pp.; Chapter 2: 140. 19
- [4] GROTHENDIECK, A.: *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV*. Inst. Hautes Études Sci. Publ. Math. **32** (1967), 361. 16
- [5] HEINS, M.: *A Holomorphic perspective of Strict Deformation Quantization*. PhD thesis, Julius-Maximilians-Universität Würzburg, October 2024. 29
- [6] JARCHOW, H.: *Locally Convex Spaces*. B. G. Teubner, Stuttgart, 1981. 1, 18, 29
- [7] KÖTHE, G.: *Topological Vector Spaces I. Grundlehren der mathematischen Wissenschaft* no. **159**. Springer-Verlag, Heidelberg, Berlin, New York, 1969. 1
- [8] KÖTHE, G.: *Topological Vector Spaces II. Grundlehren der mathematischen Wissenschaft* no. **237**. Springer-Verlag, Heidelberg, Berlin, New York, 1979. 1, 18
- [9] MCCONNELL, J. C., ROBSON, J. C.: *Noncommutative Noetherian rings*, vol. 30 in *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, revised. edition, 2001. With the cooperation of L. W. Small. 16
- [10] MEISE, R., VOGT, D.: *Einführung in die Funktionalanalysis*. Vieweg-Verlag, Braunschweig, Wiesbaden, 1992. 1

- [11] OSBORNE, M. S.: *Locally convex spaces*, vol. 269 in *Graduate Texts in Mathematics*. Springer-Verlag, Cham, Heidelberg, New York, 2014. 1, 5, 14
- [12] PIETSCH, A.: *Nuclear locally convex spaces*, vol. 66 in *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer-Verlag, New York, Heidelberg, 1972. Translated from the second German edition by William H. Ruckle. 1, 29
- [13] REED, M., SIMON, B.: *Methods of modern mathematical physics I. Functional analysis*. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 2. edition, 1980. 1
- [14] RUDIN, W.: *Functional Analysis*. McGraw-Hill Book Company, New York, 2. edition, 1991. 1
- [15] SCHAEFER, H. H.: *Topological vector spaces*, vol. 3 in *Graduate Texts in Mathematics*. Springer-Verlag, Heidelberg, New York, 2. edition, 1999. With M. P. Wolf. 1
- [16] TRÈVES, F.: *Topological vector spaces, distributions and kernels*. Dover Publications Inc., Mineola, 2006. Unabridged republication of the 1967 original. 1, 23, 26
- [17] VOGT, D.: *Lectures on Fréchet spaces*. <http://www2.math.uni-wuppertal.de/~vogt/vorlesungen/fs.pdf>, Wuppertal, Nordrhein-Westfalen, 2000. 1, 18, 29
- [18] WALDMANN, S.: *Convergence of Star Products: From Examples to a General Framework*. EMS Surv. Math. Sci. **6** (2019), 1–31. 26
- [19] WALDMANN, S.: *Differential Geometry*, Last update 2021. Unpublished lecture notes, available upon request. Currently ca. 1400 pages. 1
- [20] WALDMANN, S.: *Locally Convex Analysis*, Last update 2021. Unpublished lecture notes, available upon request. Currently ca. 300 pages. 1